

Product datasheet for **TP306615L**

Cyclin A1 (CCNA1) (NM_003914) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human cyclin A1 (CCNA1), transcript variant 1, 1 mg

Species: Human

Expression Host: HEK293T

Expression cDNA Clone or AA Sequence: >RC206615 protein sequence
Red=Cloning site **Green**=Tags(s)

METGFPAIMYPGSFIGGWGEEYLSWEGPGLPDFVFQQPVESEAMHCSNPKSGVVLATVARGPDACQILTR
APLGQDPPQRTVLGLLTANGQYRRTCGQGITRIRCYSGSENAFPPAGKKALPDCGVQEPKQGFDIYMDE
LEQGDRDSCSVREGMAFEDVYEVDTGTLKSDLHFLDFNTVSPMLVDSSLLSQSEDISSLGTDVINVTEY
AEEIYQYLREAEIRHRPKAHYMKKQPDITEGMRTILVDWLVEVGEEYKLAETLYLAVNFLDRFLSCMSV
LRGKLQLVGTAAMLLASKYEEIYPPEVDEFVYITDDTYTKRQLLMEHLLKVLAFDLTVPPTTNQFLLQY
LRRQGVCVRTENLAKYVAELSLLEADPFLKYLP SLIAAAAFCLANYTVNKHFWPETLAAFTGYSLSEIVP
CLSELHKAYLDIPHRPQQAIREKYKASKYLCVSLMEPPAVLLLQ

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Predicted MW: 52.2 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: [NP_003905](#)



[View online »](#)


Locus ID:	8900
UniProt ID:	P78396
RefSeq Size:	1965
Cytogenetics:	13q13.3
RefSeq ORF:	1509
Synonyms:	CT146

Summary: The protein encoded by this gene belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance through the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. The cyclin encoded by this gene was shown to be expressed in testis and brain, as well as in several leukemic cell lines, and is thought to primarily function in the control of the germline meiotic cell cycle. This cyclin binds both CDK2 and CDC2 kinases, which give two distinct kinase activities, one appearing in S phase, the other in G2, and thus regulate separate functions in cell cycle. This cyclin was found to bind to important cell cycle regulators, such as Rb family proteins, transcription factor E2F-1, and the p21 family proteins. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]

Protein Families: Druggable Genome

Protein Pathways: Acute myeloid leukemia, Cell cycle, Pathways in cancer, Progesterone-mediated oocyte maturation

Product images:



Coomassie blue staining of purified CCNA1 protein (Cat# [TP306615]). The protein was produced from HEK293T cells transfected with CCNA1 cDNA clone (Cat# [RC206615]) using MegaTran 2.0 (Cat# [TT210002]).