

Product datasheet for TP306638

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

RAI2 (NM_021785) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human retinoic acid induced 2 (RAI2), 20 μg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC206638 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MDDLQSQNLSMDMTDSPPALANNRLENGMAQLITTEAWNINSTDLVKKALVTVPAPSILNPPAESQSGMA LKVAATVLQPLCLGESPVVMPIHMQVEGSSAPELNPNGNATYVMTTQGPVQLPVVLEQHVFQHLNSPLVL PQEAPCSSSTIHNNLFQGAEDPEAQPQLLDLRIPSQPQEPTLPFEAVLQNLFPSQGTLGPPPCQPPPGYA PVPPQPFSSPLSPLVPPATLLVPYPVIVPLPVPVPIPIPIPVPQSSESKFSSSFPKPPSSFGLHPFKGTQ TPLEKDELKPFDILQPKEYFQLSRHTVIKMGSENEALDLSMKSVPWLKAGEVSPPIFQEDAPLDLSVAAH RKSEPPPETLYDSGASVDSSGHTVMEKLPSGMEISFAPATSHEAPAMMDSHISSSDAATEMLSQPNHPSG EVKAENNIEMVGESQAAKVIVSVEDAVPTIFCGKIKGLSGVSTKNFSFKREDSVLQGYDINSQGEESMGN

AEPLRKPIKNRSIKLKKVNSQEIHMLPIKKQRLATFFPRK

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Predicted MW: 57 kDa

Concentration: $>0.05 \mu g/\mu L$ as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by conventional

chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.





RefSeq: NP 068557

Locus ID: 10742

UniProt ID: <u>Q9Y5P3</u>, <u>A0A024RBZ8</u>, <u>B3KPD7</u>, <u>B2RBE9</u>


RefSeq Size: 2229
Cytogenetics: Xp22.13
RefSeq ORF: 1590

Summary: Retinoic acid plays a critical role in development, cellular growth, and differentiation. The

specific function of this retinoic acid-induced gene has not yet been determined but it may play a role in development. The chromosomal location of this gene designates it to be a candidate for diseases such as Nance-Horan syndrome, sensorineural deafness, non-specific X-linked cognitive disability, oral-facial-digital syndrome, and Fried syndrome. Alternate splicing results

in multiple transcript variants. [provided by RefSeq, Feb 2010]

Product images:



Coomassie blue staining of purified RAI2 protein (Cat# TP306638). The protein was produced from HEK293T cells transfected with RAI2 cDNA clone (Cat# [RC206638]) using MegaTran 2.0 (Cat# [TT210002]).