

## **Product datasheet for TP306748M**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

CN: techsupport@origene.cn

## E3 SUMO protein ligase PIAS4 (PIAS4) (NM\_015897) Human Recombinant Protein

### **Product data:**

**Product Type:** Recombinant Proteins

**Description:** Recombinant protein of human protein inhibitor of activated STAT, 4 (PIAS4), 100 μg

Species: Human
Expression Host: HEK293T

**Expression cDNA Clone** >RC206748 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MAAELVEAKNMVMSFRVSDLQMLLGFVGRSKSGLKHELVTRALQLVQFDCSPELFKKIKELYETRYAKKN SEPAPQPHRPLDPLTMHSTYDRAGAVPRTPLAGPNIDYPVLYGKYLNGLGRLPAKTLKPEVRLVKLPFFN MLDELLKPTELVPQNNEKLQESPCIFALTPRQVELIRNSRELQPGVKAVQVVLRICYSDTSCPQEDQYPP NIAVKVNHSYCSVPGYYPSNKPGVEPKRPCRPINLTHLMYLSSATNRITVTWGNYGKSYSVALYLVRQLT SSELLQRLKTIGVKHPELCKALVKEKLRLDPDSEIATTGVRVSLICPLVKMRLSVPCRAETCAHLQCFDA VFYLQMNEKKPTWMCPVCDKPAPYDQLIIDGLLSKILSECEDADEIEYLVDGSWCPIRAEKERSCSPQGA ILVLGPSDANGLLPAPSVNGSGALGSTGGGGPVGSMENGKPGADVVDLTLDSSSSSEDEEEEEEEEDED

EEGPRPKRRCPFQKGLVPAC

**TRTRPL**EQKLISEEDLAANDILDYKDDDDK**V** 

Tag: C-Myc/DDK
Predicted MW: 56.3 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

**Preparation:** Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

**Note:** For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.





**RefSeq:** NP 056981

**Locus ID:** 51588

UniProt ID: Q8N2W9, B3KMR4

RefSeq Size: 1835 Cytogenetics: 19p13.3 RefSeq ORF: 1530

Synonyms: PIAS-gamma; Piasg; PIASY; ZMIZ6

**Summary:** Functions as an E3-type small ubiquitin-like modifier (SUMO) ligase, stabilizing the interaction

between UBE2I and the substrate, and as a SUMO-tethering factor. Plays a crucial role as a transcriptional coregulation in various cellular pathways, including the STAT pathway, the p53/TP53 pathway, the Wnt pathway and the steroid hormone signaling pathway. Involved in gene silencing. Mediates sumoylation of CEBPA, PARK7, HERC2, MYB, TCF4 and RNF168. In Wnt signaling, represses LEF1 and enhances TCF4 transcriptional activities through promoting their sumoylations. Enhances the sumoylation of MTA1 and may participate in its paralog-


selective sumoylation.[UniProtKB/Swiss-Prot Function]

**Protein Families:** Transcription Factors

**Protein Pathways:** Jak-STAT signaling pathway, Pathways in cancer, Small cell lung cancer, Ubiquitin mediated

proteolysis

# **Product images:**



Coomassie blue staining of purified PIAS4 protein (Cat# [TP306748]). The protein was produced from HEK293T cells transfected with PIAS4 cDNA clone (Cat# [RC206748]) using MegaTran 2.0 (Cat# [TT210002]).