

Product datasheet for TP307216M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

JNK3 (MAPK10) (NM_002753) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human mitogen-activated protein kinase 10 (MAPK10), transcript

variant 1, 100 µg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC207216 representing NM_002753

or AA Sequence: Red=Cloning site Green=Tags(s)

MSLHFLYYCSEPTLDVKIAFCQGFDKQVDVSYIAKHYNMSKSKVDNQFYSVEVGDSTFTVLKRYQNLKPI GSGAQGIVCAAYDAVLDRNVAIKKLSRPFQNQTHAKRAYRELVLMKCVNHKNIISLLNVFTPQKTLEEFQ DVYLVMELMDANLCQVIQMELDHERMSYLLYQMLCGIKHLHSAGIIHRDLKPSNIVVKSDCTLKILDFGL ARTAGTSFMMTPYVVTRYYRAPEVILGMGYKENVDIWSVGCIMGEMVRHKILFPGRDYIDQWNKVIEQLG TPCPEFMKKLQPTVRNYVENRPKYAGLTFPKLFPDSLFPADSEHNKLKASQARDLLSKMLVIDPAKRISV DDALQHPYINVWYDPAEVEAPPPQIYDKQLDEREHTIEEWKELIYKEVMNSEEKTKNGVVKGQPSPSAQV

QQ

TRTRPLEQKLISEEDLAANDILDYKDDDDK**V**

Tag: C-Myc/DDK
Predicted MW: 48.4 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.





RefSeq: NP 002744

 Locus ID:
 5602

 UniProt ID:
 P53779

 RefSeq Size:
 2372

 Cytogenetics:
 4q21.3

 RefSeq ORF:
 1266

Synonyms: JNK3; JNK3A; p54bSAPK; p493F12; PRKM10; SAPK1b

Summary: The protein encoded by this gene is a member of the MAP kinase family. MAP kinases act as

integration points for multiple biochemical signals, and thus are involved in a wide variety of cellular processes, such as proliferation, differentiation, transcription regulation and development. This kinase is specifically expressed in a subset of neurons in the nervous system, and is activated by threonine and tyrosine phosphorylation. Targeted deletion of this

gene in mice suggests that it may have a role in stress-induced neuronal apoptosis.

Alternatively spliced transcript variants encoding different isoforms have been described for this gene. A recent study provided evidence for translational readthrough in this gene, and expression of an additional C-terminally extended isoform via the use of an alternative in-

frame translation termination codon. [provided by RefSeq, Dec 2017]


Protein Families: Druggable Genome, Protein Kinase

Protein Pathways: Adipocytokine signaling pathway, Colorectal cancer, Epithelial cell signaling in Helicobacter

pylori infection, ErbB signaling pathway, Fc epsilon RI signaling pathway, Focal adhesion, GnRH signaling pathway, Insulin signaling pathway, MAPK signaling pathway, Neurotrophin signaling pathway, NOD-like receptor signaling pathway, Pancreatic cancer, Pathways in cancer, Progesterone-mediated oocyte maturation, RIG-I-like receptor signaling pathway, Toll-like

receptor signaling pathway, Type II diabetes mellitus, Wnt signaling pathway

Product images:



Coomassie blue staining of purified MAPK10 protein (Cat# [TP307216]). The protein was produced from HEK293T cells transfected with MAPK10 cDNA clone (Cat# [RC207216]) using MegaTran 2.0 (Cat# [TT210002]).