

Product datasheet for **TP307315M**

SIRP alpha (SIRPA) (NM_080792) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human signal-regulatory protein alpha (SIRPA), transcript variant 3, 100 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA	>RC207315 protein sequence
Clone or AA Sequence:	Red=Cloning site Green=Tags(s)

MEPAGPAPGRLGPLLCLLLAASCAWSGVAGEEELQVIQPDKSVSVAAGESAILHCTVTSLIPVGPQWFR
GAGPARELIYNQKEGHFPRVTTVSESTKRENMDFSISISNITPADAGTYCVKFRKSPDTEFKSGAGTE
LSVRAKPSAPVVSFGPAARATPQHTVSFTCESHGFSRPDITLKWFKNGNELSDFQTNVDPVGESVSYSIHS
TAKVLTREDVHSQVICEVAHVTLQGDPLRGTANLSETIRVPPTLEVTQQPVRAENQVNVTCQVRKFYPQ
RLQLTWLENGNVSRTETASTVTENKDGTYNWMSWLLVNVSAHRDDVKLTCQVEHDGQPAVSKSHDLKVS
HPKEQGSNTAAENTGSNERNIYVGVVCTLLVALLMAALYLVRIRQKKAQGSTSSTRLHEPEKNAREIT
QDNDITYADLNLPGKPKPAPQAAEPNNHTEYASIQTSPQASEDTLTYADLDMVHLNRTPKQPAPKPEP
SFSEYASVQVPRK

TRTRPLEQKLISEEDLAANDILDYKDDDDKV


Tag:	C-Myc/DDK
Predicted MW:	52.3 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.



[View online »](#)

Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_542970
Locus ID:	140885
UniProt ID:	P78324
RefSeq Size:	3868
Cytogenetics:	20p13
RefSeq ORF:	1509
Synonyms:	BIT; CD172A; MFR; MYD-1; P84; PTPNS1; SHPS1; SIRP
Summary:	The protein encoded by this gene is a member of the signal-regulatory-protein (SIRP) family, and also belongs to the immunoglobulin superfamily. SIRP family members are receptor-type transmembrane glycoproteins known to be involved in the negative regulation of receptor tyrosine kinase-coupled signaling processes. This protein can be phosphorylated by tyrosine kinases. The phospho-tyrosine residues of this PTP have been shown to recruit SH2 domain containing tyrosine phosphatases (PTP), and serve as substrates of PTPs. This protein was found to participate in signal transduction mediated by various growth factor receptors. CD47 has been demonstrated to be a ligand for this receptor protein. This gene and its product share very high similarity with several other members of the SIRP family. These related genes are located in close proximity to each other on chromosome 20p13. Multiple alternatively spliced transcript variants have been determined for this gene. [provided by RefSeq, Jul 2008]
Protein Families:	Druggable Genome, Phosphatase, Transmembrane

Product images:



Coomassie blue staining of purified SIRPA protein (Cat# [TP307315]). The protein was produced from HEK293T cells transfected with SIRPA cDNA clone (Cat# [RC207315]) using MegaTran 2.0 (Cat# [TT210002]).