

Product datasheet for TP307333

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

EIF4E (NM_001968) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human eukaryotic translation initiation factor 4E (EIF4E), transcript

variant 1, 20 µg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC207333 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MATVEPETTPTPNPPTTEEEKTESNQEVANPEHYIKHPLQNRWALWFFKNDKSKTWQANLRLISKFDTVE DFWALYNHIQLSSNLMPGCDYSLFKDGIEPMWEDEKNKRGGRWLITLNKQQRRSDLDRFWLETLLCLIGE SFDDYSDDVCGAVVNVRAKGDKIAIWTTECENREAVTHIGRVYKERLGLPPKIVIGYQSHADTATKSGST

TKNRFVV

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Predicted MW: 24.9 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 001959

Locus ID: 1977





UniProt ID: P06730 4749 RefSeq Size: Cytogenetics: 4q23 RefSeq ORF: 651

Synonyms: AUTS19; CBP; eIF-4E; EIF4E1; EIF4EL1; EIF4F

Summary: The protein encoded by this gene is a component of the eukaryotic translation initiation factor


4F complex, which recognizes the 7-methylguanosine cap structure at the 5' end of

messenger RNAs. The encoded protein aids in translation initiation by recruiting ribosomes to the 5'-cap structure. Association of this protein with the 4F complex is the rate-limiting step in translation initiation. This gene acts as a proto-oncogene, and its expression and activation is associated with transformation and tumorigenesis. Several pseudogenes of this gene are found on other chromosomes. Alternative splicing results in multiple transcript variants.

[provided by RefSeq, Sep 2015]

Protein Pathways: Insulin signaling pathway, mTOR signaling pathway

Product images:



Coomassie blue staining of purified EIF4E protein (Cat# TP307333). The protein was produced from HEK293T cells transfected with EIF4E cDNA clone (Cat# [RC207333]) using MegaTran 2.0 (Cat# [TT210002]).