

Product datasheet for TP307354L

PABP (PABPC1) (NM_002568) Human Recombinant Protein

Product data:


Product Type:	Recombinant Proteins
Description:	Recombinant protein of human poly(A) binding protein, cytoplasmic 1 (PABPC1), 1 mg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC207354 representing NM_002568 Red=Cloning site Green=Tags(s)
	<p>MNPSAPSYPMASLYVGDLPDVTEAMLYEKFSPAGPILSIRVCRDMITRRSLGYAYVNFQQPADAERALD TMNFDVIKGPVVRIMWSQRDPSLRKSGVGNIFIKNLDKSIDNKALYDTFSAFGNILSCKWVCDENGSKGY GFVHFETQEAERAIEKMNGMLLNDRKVFVGRFKSRKEREALGARAKEFTNVYIKNFGEDMDDERLKD LFGKFGPALSVMKMTDESCKSGFGFVSFERHEDAQKAVDEMNGKELNGKQIYVGRAQKKVERQTELKRKF EQMKQDRITRYQGVNLYVKNLDDGIDDERLRKEFSPFGTITSAKVMMEGGRSKGFGFVCFSSPEEATKAV TEMNGRIVATKPLYVALAQRKEERQAHLTNQYMQRMASVRAVNPVINPYQPAPPSGYFMAAIPQTQNR AAYPPSQIAQLRPSRWTAQGARPHPFQNMPGAIRPAAPRPPFSTMRPASSQVPRVMSTQRVANTSTQTM GPRPAAAAAATPAVRTVPQYKYAAGVRNPQQHLNAQPQVTMQQPAVHVQGQEPLTASMLASAPPQEQKQ MLGERLFLPIQAMHPTLAGKITGMILLEIDNSELLHMLESPESLRSKVDEAVAVLQAHQAKEAAQKAVNSA TGVPTV</p> <p>SGPTRRRLEQKLISEEDLAANDILDYKDDDDKV</p>
Tag:	C-Myc/DDK
Predicted MW:	70.5 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.



[View online »](#)

Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	<u>NP_002559</u>
Locus ID:	26986
UniProt ID:	<u>P11940, A0A024R9C1</u>
RefSeq Size:	2869
Cytogenetics:	8q22.3
RefSeq ORF:	1908
Synonyms:	PAB1; PABP; PABP1; PABPC2; PABPL1
Summary:	This gene encodes a poly(A) binding protein. The protein shuttles between the nucleus and cytoplasm and binds to the 3' poly(A) tail of eukaryotic messenger RNAs via RNA-recognition motifs. The binding of this protein to poly(A) promotes ribosome recruitment and translation initiation; it is also required for poly(A) shortening which is the first step in mRNA decay. The gene is part of a small gene family including three protein-coding genes and several pseudogenes. [provided by RefSeq, Aug 2010]

Product images:



Coomassie blue staining of purified PABPC1 protein (Cat# [TP307354]). The protein was produced from HEK293T cells transfected with PABPC1 cDNA clone (Cat# [RC207354]) using MegaTran 2.0 (Cat# [TT210002]).