

Product datasheet for TP307506L

DIP13B (APPL2) (NM_018171) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human adaptor protein, phosphotyrosine interaction, PH domain and leucine zipper containing 2 (APPL2), 1 mg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC207506 protein sequence Red=Cloning site Green=Tags(s)

MPAVDKLLLEALQDSPQTRSLLSVFEEDAGTLTDYTNQLLQAMQRVYGAQNEMCLATQQLSKQLLAYEK
QNFALGKGDDEEVISTLHYFSKWDELNLLHTELAQLADTMVLPPIIQFREKDLTEVSTLKDLFGLASNEH
DLSMAKYSRLPKKKENEKVKTEVGKEVAAARRKQHLSSLQYYCALNALQYRKQMAMMEPMIGFAHGQINF
FKKGAEMFSKRMDNFLSSVADMVQSIQVELEAEAEKMRVSSQELLSVDESUYTPDSDVAAPQINRNLQK
AGYLNLRNKTGLVTTTWERLYFFTQGGNLMCQPRGAVAGGLIQDLNCSVMAVDCEDRRYCFQITTPNGK
SGIILQAESRKENEWICAINNISRQIYLTDNPEAVAIKLNQTALQAVTPITSEFGKKQESSCPSQNLKNS
EMENENDKIVPKVTASLPEAEELIAPGTPIQFDIVLPATEFLDQNRGSRRTNPFGETEDESFPEDSLL
QQMFIVRFLGSMVAVKTDSTTEVIYEAMRQVLAARAIHNIFRMTESHLMVTSQSLRLIDPQTQVSRANFEL
TSVTQFAAHQENKRLVGFVIRVPESTGEESLSTYIFESNSEGEKICYAINLGKEIIEVQKDPEALQMLL
SIPLTNDGKYVLLNDQPDDDDGNPNEHRAESEA

TRTRPLEQKLISEEDLAANDILDYKDDDDKV


Tag:	C-Myc/DDK
Predicted MW:	74.3 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.



[View online »](#)

Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_060641
Locus ID:	55198
UniProt ID:	Q8NEU8
RefSeq Size:	3289
Cytogenetics:	12q23.3
RefSeq ORF:	1992
Synonyms:	DIP13B
Summary:	The protein encoded by this gene is one of two effectors of the small GTPase RAB5A/Rab5, which are involved in a signal transduction pathway. Both effectors contain an N-terminal Bin/Amphiphysin/Rvs (BAR) domain, a central pleckstrin homology (PH) domain, and a C-terminal phosphotyrosine binding (PTB) domain, and they bind the Rab5 through the BAR domain. They are associated with endosomal membranes and can be translocated to the nucleus in response to the EGF stimulus. They interact with the NuRD/MeCP1 complex (nucleosome remodeling and deacetylase /methyl-CpG-binding protein 1 complex) and are required for efficient cell proliferation. A chromosomal aberration t(12;22)(q24.1;q13.3) involving this gene and the PSAP2 gene results in 22q13.3 deletion syndrome, also known as Phelan-McDermid syndrome. [provided by RefSeq, Oct 2011]

Product images:



Coomassie blue staining of purified APPL2 protein (Cat# [TP307506]). The protein was produced from HEK293T cells transfected with APPL2 cDNA clone (Cat# [RC207506]) using MegaTran 2.0 (Cat# [TT210002]).