

Product datasheet for TP307809

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PRKAR1B (NM_002735) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human protein kinase, cAMP-dependent, regulatory, type I, beta

(PRKAR1B), 20 µg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC207809 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MASPPACPSEEDESLKGCELYVQLHGIQQVLKDCIVHLCISKPERPMKFLREHFEKLEKEENRQILARQK SNSQSDSHDEEVSPTPPNPVVKARRRRGGVSAEVYTEEDAVSYVRKVIPKDYKTMTALAKAISKNVLFAH LDDNERSDIFDAMFPVTHIAGETVIQQGNEGDNFYVVDQGEVDVYVNGEWVTNISEGGSFGELALIYGTP RAATVKAKTDLKLWGIDRDSYRRILMGSTLRKRKMYEEFLSKVSILESLEKWERLTVADALEPVQFEDGE KIVVQGEPGDDFYIITEGTASVLQRRSPNEEYVEVGRLGPSDYFGEIALLLNRPRAATVVARGPLKCVKL

DRPRFERVLGPCSEILKRNIQRYNSFISLTV

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK
Predicted MW: 42.9 kDa

Concentration: >0.05 μg/μL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 002726





Locus ID: 5575

UniProt ID: P31321
RefSeq Size: 2512
Cytogenetics: 7p22.3
RefSeq ORF: 1143
Synonyms: PRKAR1

Summary: The protein encoded by this gene is a regulatory subunit of cyclic AMP-dependent protein


kinase A (PKA), which is involved in the signaling pathway of the second messenger cAMP. Two regulatory and two catalytic subunits form the PKA holoenzyme, disbands after cAMP binding. The holoenzyme is involved in many cellular events, including ion transport, metabolism, and transcription. Several transcript variants encoding the same protein have

been found for this gene. [provided by RefSeq, Aug 2015]

Protein Families: Druggable Genome

Protein Pathways: Apoptosis, Insulin signaling pathway

Product images:



Coomassie blue staining of purified PRKAR1B protein (Cat# TP307809). The protein was produced from HEK293T cells transfected with PRKAR1B cDNA clone (Cat# [RC207809]) using MegaTran 2.0 (Cat# [TT210002]).