

Product datasheet for **TP307976L**

Solo (SESTD1) (NM_178123) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human SEC14 and spectrin domains 1 (SESTD1), 1 mg
Species:	Human
Expression Host:	HEK293T
Expression cDNA	>RC207976 protein sequence
Clone or AA Sequence:	Red=Cloning site Green=Tags(s)

MEASVILPILKKKLAFLSGGKDRRSLILTIPLCLEQTNMDELSVTLDFLLSIPSEKCKARGFTVIVDGR
KSQWNVKTWVWMLQNWVPAEVSVCVWKPDEFWDKVKVTHFCFWKEKDRLGFEVILVSANKLTRYIEPCQ
LTEDFGGSLTYDHMDWLNKRLVFEKFTKESTSLLELALINNGSDKGNQKEKERSVDLNFPSVDPETVL
QTGHELLSELQRRFNGSDGGVSWSPMDDELLAQPQVMKLLDSLREQYTRYQEVCRQRSKRTQLEEIQQK
VMQVNWLEGPGEQLRAQWIGDSIRASQALQQKHEEIESQHSEWFVAVYVELNQIAALLNAGDEEDLV
ELKSLQQQLSDVCYRQASQLEFRQNLQAALFHVGAQDLSQQLDGLLGMLCVDVAPADGASIQQTLKLL
EEKLKSVDVGLQGLREKGGQLLDQISNQASWAYGKDVTIENKENVVDHIQGVMEQMLRQRCEMMDVRR
LKMLQMVQLFKCEEDAAQAVEWSELDDALLKTHIRLGDDAQETKVLEKHRKFVDVAQSTYDYGRQLLQ
ATVWLCQSLRCTSRSSGDTLPRNLNRVWKQFTIASEERVHRLEMAIAFHSAEKILQDCPEEPEAINDEEQ
FDEIEAVGKSLDRLTVPVVPDGTQYFGSPSDMASTAENIRDRMKLVNLKRQQLRHPPEMVTES

TRTRPLEQKLISEEDLAANDILDYKDDDDKV


Tag:	C-Myc/DDK
Predicted MW:	79.2 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.



[View online »](#)

Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_835224
Locus ID:	91404
UniProt ID:	Q86VW0 , B3KTX3
RefSeq Size:	10448
Cytogenetics:	2q31.2
RefSeq ORF:	2088
Synonyms:	SOLO
Summary:	May act as the primary docking protein directing membrane turnover and assembly of the transient receptor potential channels TRPC4 and TRPC5. Binds phospholipids such as phosphatidylinositol monophosphates, phosphatidylinositol diphosphates (PIP2s) and phosphatidic acid, but not less polar lipids including phosphatidylcholine, phosphatidylserine, and phosphatidylinositol. The binding to PIP2s is calcium dependent. Might be involved in the plasma membrane localization of CTNNB1.[UniProtKB/Swiss-Prot Function]
Protein Families:	Druggable Genome

Product images:



Coomassie blue staining of purified SESTD1 protein (Cat# [TP307976]). The protein was produced from HEK293T cells transfected with SESTD1 cDNA clone (Cat# [RC207976]) using MegaTran 2.0 (Cat# [TT210002]).