

Product datasheet for **TP308198**

PDP1 (NM_018444) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins
Description: Recombinant protein of human protein phosphatase 2C, magnesium-dependent, catalytic subunit (PPM2C), nuclear gene encoding mitochondrial protein, 20 µg
Species: Human
Expression Host: HEK293T
Expression cDNA >RC208198 protein sequence
Clone or AA Sequence: Red=Cloning site Green=Tags(s)

MPAPTQLFFPLIRNCELSRIYGTACYCHHKHLCCSSSYIPQSRLRYTPHPAYATFCRPKENWWQYTQGRR
YASTPQKFYLTTPQVNSILKANEYSFKVPEFDGKNVSSILGFDSNQLPANAPIEDRRSAATCLQTRGMLL
GVFDGHAGCACSQAVSERLFYYIAVSLPHETLLEIENAVESGRALLPILQWHKHPNDYFSKEASKLYFN
SLRTYWQELIDLNTGESTDIDVKEALINAFKRLDNDISLEAQVGDPNFLNYLVLRVAFSGATACVAHVD
GVDLHVANTGDSRAMLGVQEEDGSWSAVTSLNDHNAQNERELERLKHHPKSEAKSVKQDRLLGLLMPF
RAFGDVKFKWSIDLQKRVIESGPDQLNDNEYTKFIPPNYHTPPYLTAPEVYHRLRPQDKFLVLTADGL
WETMHRQDWRIVGEYLTGMHHQQPIAVGGYKVTLGQMHGLLTERRTKMSSVFEDQNAATHLIRHAVGNN
EFGTVDHERLSKMLSLPEELARMYRDDITIIVVQFNSHVVGAYQNQE

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK
Predicted MW: 60.9 kDa
Concentration: >0.05 µg/µL as determined by microplate BCA method
Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation: Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note: For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage: Store at -80°C.




[View online »](#)

Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_060914
Locus ID:	54704
UniProt ID:	Q9P0J1 , A0A024R9C0
RefSeq Size:	4291
Cytogenetics:	8q22.1
RefSeq ORF:	1611
Synonyms:	PDH; PDP; PDPC; PPM2A; PPM2C
Summary:	Pyruvate dehydrogenase (E1) is one of the three components (E1, E2, and E3) of the large pyruvate dehydrogenase complex. Pyruvate dehydrogenase kinases catalyze phosphorylation of serine residues of E1 to inactivate the E1 component and inhibit the complex. Pyruvate dehydrogenase phosphatases catalyze the dephosphorylation and activation of the E1 component to reverse the effects of pyruvate dehydrogenase kinases. Pyruvate dehydrogenase phosphatase is a heterodimer consisting of catalytic and regulatory subunits. Two catalytic subunits have been reported; one is predominantly expressed in skeletal muscle and another one is much more abundant in the liver. The catalytic subunit, encoded by this gene, is the former, and belongs to the protein phosphatase 2C (PP2C) superfamily. Along with the pyruvate dehydrogenase complex and pyruvate dehydrogenase kinases, this enzyme is located in the mitochondrial matrix. Mutation in this gene causes pyruvate dehydrogenase phosphatase deficiency. Multiple alternatively spliced transcript variants encoding different isoforms have been identified.[provided by RefSeq, Jun 2009]

Protein Families: Druggable Genome, Phosphatase

Product images:



Coomassie blue staining of purified PDP1 protein (Cat# TP308198). The protein was produced from HEK293T cells transfected with PDP1 cDNA clone (Cat# [RC208198]) using MegaTran 2.0 (Cat# [TT210002]).