

Product datasheet for TP308387

Staufen (STAU1) (NM_017453) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human staufen, RNA binding protein, homolog 1 (Drosophila) (STAU1), transcript variant T3, 20 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC208387 representing NM_017453 Red =Cloning site Green =Tags(s)

MSQVQVQVQNPSAALSGSQILNKNQSLLSQPLMSIPSTTSSLPSENAGRPIQNSALPSASITSTSAAAES
ITPTVELNALCMKLGKKPMYKVPDPYSRMQSTYNYNMRGGAYPPRYFYPFPVPLLYQVELSVGGQQFNG
K GKTRQAAKHDAAAKALRILQNEPLPERLEVNGRESEENLNKSEISQVFEIALKRNLVNFVARESGP
PHMKNFVTKVSVGEFVGE GEGKSKKISKNAAIAVLEELKKLPPLPAVERVKPRIKKKTKPIVKPQTSPE
YGQGINPISRLAQIQAKKEKEPEYLLTERGLPRRREFVMQVKVGNHTAEGTGTNKKVAKRNAENMLE
ILGFKVPQAQPTKPALKSEEKTPIKKPGDGRKVTFFEPGSGDENGTSNKEDEFRMPYLSHQQLPAGILPM
VPEVAQAVGVSQGHHTKDFTRAAPNPAKATVTAMIARELLYGGTSPTAETILKNNISSGHVPHGPLTRPS
EQLDYLRSRVQGFQVEYKDFPKNNKNEFVSLINCSSQPPLISHGIGKDVESCHDMAALNILKLLSELDQQS
TEMPRTGNGPMSVCGRC

TRTRPLEQKLISEEDLAANDILDYKDDDDKV


Tag:	C-Myc/DDK
Predicted MW:	63 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.



[View online »](#)

Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_059347
Locus ID:	6780
UniProt ID:	O95793 , B3KRE0
RefSeq Size:	3688
Cytogenetics:	20q13.13
RefSeq ORF:	1731
Synonyms:	PPP1R150; STAU
Summary:	Staufen is a member of the family of double-stranded RNA (dsRNA)-binding proteins involved in the transport and/or localization of mRNAs to different subcellular compartments and/or organelles. These proteins are characterized by the presence of multiple dsRNA-binding domains which are required to bind RNAs having double-stranded secondary structures. The human homologue of staufen encoded by STAU, in addition contains a microtubule-binding domain similar to that of microtubule-associated protein 1B, and binds tubulin. The STAU gene product has been shown to be present in the cytoplasm in association with the rough endoplasmic reticulum (RER), implicating this protein in the transport of mRNA via the microtubule network to the RER, the site of translation. [provided by RefSeq, Apr 2020]

Product images:



Coomassie blue staining of purified STAU1 protein (Cat# TP308387). The protein was produced from HEK293T cells transfected with STAU1 cDNA clone (Cat# [RC208387]) using MegaTran 2.0 (Cat# [TT210002]).