

Product datasheet for **TP308397M**

AMH (NM_000479) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human anti-Mullerian hormone (AMH), 100 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC208397 representing NM_000479 Red =Cloning site Green =Tags(s)

MRDLPLTSLALVLSALGALLGTEALRAEEPAVGTSGLI FREDLDWPPGSPQEPLCLVALGGDSNGSSSPL
RVGALSAYEQAFLGAVQRRARWGPRDLATFGVCNTGDRQAALPSLRRLGAWLRDPGGQRLVHLHEEVTW
EPTPSLRFQEPGGAGPPELALLVLYPGPGPEVTVTRAGLPGAQSLCPSRDTRYLVLAVDRPAGAWRGS
GLALTLQPRGEDSRLSTARLQALLFGDDHRCFTRMTPALLLLPRSEAPLPAHGQLDTPFPFPPRPSAEL
EESPPSADPFLETLTRLVLRALRVPPARASAPRLALDPDALAGFPQGLVNLSDPAALERLLDGEEPLLLLL
RPTAATTGDPAPLHDPTSAPWATALARRVAAELQAAAAELRSLPGLPPATAPLLARLLALCPGGPGGLGD
PLRALLLLKALQGLRVEWRGRDPRGPGRAQRSAGATAADGPCALRELSVDLRAERSVLIPETYQANNCQG
VCGWPQSDRNPRYGNHVLLLLKMQARGAALARPPCCVPTAYAGKLLISLSEERISAHHVPMNVATECGCR

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag:	C-Myc/DDK
Predicted MW:	56.6 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Bioactivity:	In vivo treatment (PMID: 29760445)
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.



[View online »](#)

Stability: Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: [NP_000470](#)

Locus ID: 268

UniProt ID: [P03971](#)

RefSeq Size: 2008

Cytogenetics: 19p13.3

RefSeq ORF: 1680


Synonyms: MIF; MIS

Summary: This gene encodes a secreted ligand of the TGF-beta (transforming growth factor-beta) superfamily of proteins. Ligands of this family bind various TGF-beta receptors leading to recruitment and activation of SMAD family transcription factors that regulate gene expression. The encoded preproprotein is proteolytically processed to generate N- and C-terminal cleavage products that homodimerize and associate to form a biologically active noncovalent complex. This complex binds to the anti-Mullerian hormone receptor type 2 and causes the regression of Mullerian ducts in the male embryo that would otherwise differentiate into the uterus and fallopian tubes. This protein also plays a role in Leydig cell differentiation and function and follicular development in adult females. Mutations in this gene result in persistent Mullerian duct syndrome. [provided by RefSeq, Jul 2016]

Protein Families: Druggable Genome, Secreted Protein

Protein Pathways: Cytokine-cytokine receptor interaction, TGF-beta signaling pathway

Product images:



Coomassie blue staining of purified AMH protein (Cat# [TP308397]). The protein was produced from HEK293T cells transfected with AMH cDNA clone (Cat# [RC208397]) using MegaTran 2.0 (Cat# [TT210002]).