

Product datasheet for TP308522L

TAB1 (NM_006116) Human Recombinant Protein

Product data:


Product Type:	Recombinant Proteins
Description:	Recombinant protein of human mitogen-activated protein kinase kinase kinase 7 interacting protein 1 (MAP3K7IP1), transcript variant alpha, 1 mg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC208522 protein sequence Red =Cloning site Green =Tags(s)
	<p>MAAQRRSLLQSEQQPSWTDLPLCHLSGVGSASNRSYSADGKGTESHPPEDSWLKFRSENNCFLYGVFNG YDGNRVTNFVAQRLSAELLGQLNAEHAHADVRRVLLQAFDVVERSFLSIDDALAEKASLQSQLPEGVP QHQLPPQYQKILERLKTLEISGGAMAVVAVLLNNKLYVANVGTNRALLCKSTVDGLQVTQLNVDHTTE NEDELFRLSQLGLDAGKIKQVGIICGQESTRRIGDYKVYGYTDIDLSSAAKSKPIIAEPEIHGAQPLDG VTGFLVLMSEGLYKALEAAHGPGQANQEIAAMIDTEFAKQTSLDAVAQAVVDRVKRIHSDFASGGERAR FCPRHEDMTLLVRNFGYPLGEMSQPTSPAPAAGGRVYPVSVPYSSAQSTSKTSVTLVSLVMP SQGMVNG AHSASTLDEATPLTNQSPTLTLQSTNHTQSSSSSSDGGFLRSRPAHSLPPGEDGRVEPYVDFAEFYRL WSVDHGEQSVTAP</p> <p>TRTRPLEQKLISEEDLAANDILDYKDDDDKV</p>
Tag:	C-Myc/DDK
Predicted MW:	54.5 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.



[View online »](#)

Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_006107
Locus ID:	10454
UniProt ID:	Q15750 , A8K6K3
RefSeq Size:	3240
Cytogenetics:	22q13.1
RefSeq ORF:	1512
Synonyms:	3'-Tab1; MAP3K7IP1
Summary:	The protein encoded by this gene was identified as a regulator of the MAP kinase kinase kinase MAP3K7/TAK1, which is known to mediate various intracellular signaling pathways, such as those induced by TGF beta, interleukin 1, and WNT-1. This protein interacts and thus activates TAK1 kinase. It has been shown that the C-terminal portion of this protein is sufficient for binding and activation of TAK1, while a portion of the N-terminus acts as a dominant-negative inhibitor of TGF beta, suggesting that this protein may function as a mediator between TGF beta receptors and TAK1. This protein can also interact with and activate the mitogen-activated protein kinase 14 (MAPK14/p38alpha), and thus represents an alternative activation pathway, in addition to the MAPKK pathways, which contributes to the biological responses of MAPK14 to various stimuli. Alternatively spliced transcript variants encoding distinct isoforms have been reported. [provided by RefSeq, Jul 2008]
Protein Families:	Druggable Genome
Protein Pathways:	MAPK signaling pathway, NOD-like receptor signaling pathway, Toll-like receptor signaling pathway

Product images:



Coomassie blue staining of purified TAB1 protein (Cat# [TP308522]). The protein was produced from HEK293T cells transfected with TAB1 cDNA clone (Cat# [RC208522]) using MegaTran 2.0 (Cat# [TT210002]).