

## Product datasheet for **TP308568L**

### TXNDC5 (NM\_022085) Human Recombinant Protein

#### Product data:


Product Type:	Recombinant Proteins
Description:	Recombinant protein of human thioredoxin domain containing 5 (TXNDC5), transcript variant 2, 1 mg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC208568 representing NM_022085 <b>Red</b> =Cloning site <b>Green</b> =Tags(s)
	MTQSVDSNRGNRNEKRCGHCQRLQPTWNDLGDKYNSMEDAKVYVAKVDCTAHSVCSAQGVRGYPTLKLFPKPGQEAVKYQGPRDFQTLNWMQLTLNEEPVTPEPEVEPPSAPELKQGLYELASNFELHVAQGDHFIKFFAPWCGHCKALAPTWEQLALGLEHSETVKIGKVDCTQHVELCSGNQVRGYPTLLWFRDGGKVDQYKGRDLESLREYVESQLQRTETGATETVTPSEAPVLAPEADKGTVLALTENNFDTTIAEGITFIKFIYAPWCGHCKTLAPTWEELSKKEFPGLAGVKIAEVDCTAERNICKYSVRYPTLLLFRGGKVKVSEHSGGRDLDSLHRFVLSQAKDEL
	<b>TRTRPLEQKLISEEDLAANDILDYKDDDDKV</b>
Tag:	C-Myc/DDK
Predicted MW:	43.5 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	<u><a href="#">NP_071368</a></u>



[View online »](#)

Locus ID:	81567
UniProt ID:	<a href="#">Q8NBS9</a>
RefSeq Size:	3047
Cytogenetics:	6p24.3
RefSeq ORF:	1170
Synonyms:	ERP46, Hcc-2, UNQ364, EndoPDI, MGC3178; thioredoxin domain containing 5
Summary:	This gene encodes a member of the disulfide isomerase (PDI) family of endoplasmic reticulum (ER) proteins that catalyze protein folding and thiol-disulfide interchange reactions. The encoded protein has an N-terminal endoplasmic reticulum (ER)-signal sequence, three catalytically active thioredoxin domains and a C-terminal ER-retention sequence. Its expression is induced by hypoxia and its role may be to protect hypoxic cells from apoptosis. Alternative splicing results in multiple transcript variants. Read-through transcription also exists between this gene and the neighboring upstream BLOC1S5 gene. [provided by RefSeq, Dec 2016]
Protein Families:	Druggable Genome

### Product images:



Coomassie blue staining of purified TXNDC5 protein (Cat# [TP308568]). The protein was produced from HEK293T cells transfected with TXNDC5 cDNA clone (Cat# [RC208568]) using MegaTran 2.0 (Cat# [TT210002]).