

Product datasheet for **TP308675M**

BYSL (NM_004053) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human bystin-like (BYSL), 100 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC208675 protein sequence Red =Cloning site Green =Tags(s) MPKFKAARGVGGQEKHAPLADQILAGNAVVRAGVREKRRGRGTGEAEFEYVGPRLSRRILQQARQQQEELE AEHGTGDKPAAPRERTTRLGPRMPQDGSDDDEEWPTEKAATMTAAGHHAEVVDPEDERAIEFMNKN PPARRTLADIIMEKLTEKQTEVETVMSEVSGFPMPQLDPRVLEVYRGVREVLKYSRSGKLPKAFKIIPAL SNWEQILYVTEPEAWTAAAMYQATRIFASNLERMAQRFYNLVLPRVRDDVAEYKRLNFHLYMALKKAL FKPGAWFKGILIPCESGTCTLREAIIVGSIITKCSIPVLHSSAAMLKIAEMEYSGANSIFLRLLLDKKY ALPYRVLDALVFHFLGFRTEKRELPLVWHQCLLTLVQRYKADLATDQKEALLELLRLQHPQLSPEIRRE LQSAVPRDVEDVPITVE TRTRPLEQKLISEEDLAANDILDYKDDDDKV
Tag:	C-Myc/DDK
Predicted MW:	49.4 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	<u>NP_004044</u>



[View online »](#)

Locus ID: 705

UniProt ID: [Q13895](#)

RefSeq Size: 2005

Cytogenetics: 6p21.1


RefSeq ORF: 1311

Synonyms: BYSTIN; Enp1

Summary: Bystin is expressed as a 2-kb major transcript and a 3.6-kb minor transcript in SNG-M cells and in human trophoblastic teratocarcinoma HT-H cells. Protein binding assays determined that bystin binds directly to trophinin and tastin, and that binding is enhanced when cytokeratins 8 and 18 are present. Immunocytochemistry of HT-H cells showed that bystin colocalizes with trophinin, tastin, and the cytokeratins, suggesting that these molecules form a complex in trophectoderm cells at the time of implantation. Using immunohistochemistry it was determined that trophinin and bystin are found in the placenta from the sixth week of pregnancy. Both proteins were localized in the cytoplasm of the syncytiotrophoblast in the chorionic villi and in endometrial decidual cells at the uteroplacental interface. After week 10, the levels of trophinin, tastin, and bystin decreased and then disappeared from placental villi. [provided by RefSeq, Jul 2008]

Protein Families: Stem cell - Pluripotency

Product images:



Coomassie blue staining of purified BYSL protein (Cat# [TP308675]). The protein was produced from HEK293T cells transfected with BYSL cDNA clone (Cat# [RC208675]) using MegaTran 2.0 (Cat# [TT210002]).