

Product datasheet for **TP308776**

MID1 (NM_000381) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Purified recombinant protein of Homo sapiens midline 1 (Opitz/BBB syndrome) (MID1), transcript variant 1, 20 µg

Species: Human

Expression Host: HEK293T

Expression cDNA Clone or AA Sequence: >RC208776 protein sequence
Red=Cloning site **Green**=Tags(s)

METLESELTCPICLELFEDPLLLPCAHSFCNCAHRILVSHCATNESVESITAFQCPTCRHVITLSQRGL
DGLKRNVTLQNIIDRFQKASVSGPNPSETRRERAFDANTMTSAEKVLCQFCDQDPAQDAVKTCVTCEVS
YCDECLKATHPNKKPFTGHRLEIPDISHIRGLMCLEHEDEKVNMYCVTDDQLICALCKLVGRHRDHQVA
ALSERYDKLKQNLESNLTNLIKRNTELETLLAKLIQTCQHVEVNASRQEAKLTEECDLLIEIIQRRQII
GTKIKEGKVMRLRKLAAQQIANCKQCIERSASLISQAEHSLKENDHARFLQTAKNITERVSMATASSQVLI
PEINLNDTFDFTALDFSREKKLLECLDYLTAPNPPTIREELCTASYDTITVHWTSDDFEFSVSYELQYTI
FTGQANVVSLCNSADSWMIVPNIKQNHVTVHGLQSGTKYIFMVKAINQAGSRSSSEPGKLTNSQPFKLDP
KSAHRKLVSHDNLTVRDESSKKSHTPERFTSQGSYGVAGNVFIDSGRHYWEVWISGSTWYAIGLAYK
SAPKHEWIGKNSASWALCRCNNNWWVRHNSKEIPIEPAPHLRRVGILLDYDNGSIAFYDALNSIHLYTFD
VAFAQPVCPTFTVWNKCLTIITGLPIPDHLDCTEQLP

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Predicted MW: 75.1 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.


Storage: Store at -80°C.



[View online »](#)

Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_000372
Locus ID:	4281
UniProt ID:	O15344 , A0A024RBV4
RefSeq Size:	6463
Cytogenetics:	Xp22.2
RefSeq ORF:	2001
Synonyms:	BBBG1; FXY; GBBB1; MIDIN; OGS1; OS; OSX; RNF59; TRIM18; XPRF; ZNFX
Summary:	The protein encoded by this gene is a member of the tripartite motif (TRIM) family, also known as the 'RING-B box-coiled coil' (RBCC) subgroup of RING finger proteins. The TRIM motif includes three zinc-binding domains, a RING, a B-box type 1 and a B-box type 2, and a coiled-coil region. This protein forms homodimers which associate with microtubules in the cytoplasm. The protein is likely involved in the formation of multiprotein structures acting as anchor points to microtubules. Mutations in this gene have been associated with the X-linked form of Opitz syndrome, which is characterized by midline abnormalities such as cleft lip, laryngeal cleft, heart defects, hypospadias, and agenesis of the corpus callosum. This gene was also the first example of a gene subject to X inactivation in human while escaping it in mouse. Alternative promoter use, alternative splicing and alternative polyadenylation result in multiple transcript variants that have different tissue specificities. [provided by RefSeq, Dec 2016]
Protein Families:	Druggable Genome
Protein Pathways:	Ubiquitin mediated proteolysis

Product images:



Coomassie blue staining of purified MID1 protein (Cat# TP308776). The protein was produced from HEK293T cells transfected with MID1 cDNA clone (Cat# [RC208776]) using MegaTran 2.0 (Cat# [TT210002]).