

Product datasheet for TP308975L

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

ICOS Ligand (ICOSLG) (NM_015259) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human inducible T-cell co-stimulator ligand (ICOSLG), 1 mg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC208975 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MRLGSPGLLFLLFSSLRADTQEKEVRAMVGSDVELSCACPEGSRFDLNDVYVYWQTSESKTVVTYHIPQN SSLENVDSRYRNRALMSPAGMLRGDFSLRLFNVTPQDEQKFHCLVLSQSLGFQEVLSVEVTLHVAANFSV PVVSAPHSPSQDELTFTCTSINGYPRPNVYWINKTDNSLLDQALQNDTVFLNMRGLYDVVSVLRIARTPS VNIGCCIENVLLQQNLTVGSQTGNDIGERDKITENPVSTGEKNAATWSILAVLCLLVVVAVAIGWVCRDR

CLQHSYAGAWAVSPETELTGHV

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Predicted MW: 33.2 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 056074

Locus ID: 23308





ICOS Ligand (ICOSLG) (NM_015259) Human Recombinant Protein - TP308975L

UniProt ID: <u>075144</u>, <u>A0N0L8</u>

RefSeq Size: 3320 Cytogenetics: 21q22.3 RefSeq ORF: 906

Synonyms: B7-H2; B7h; B7H2; B7RP-1; B7RP1; CD275; GL50; ICOS-L; ICOSL; LICOS

Summary: Ligand for the T-cell-specific cell surface receptor ICOS. Acts as a costimulatory signal for T-cell

proliferation and cytokine secretion; induces also B-cell proliferation and differentiation into plasma cells. Could play an important role in mediating local tissue responses to inflammatory


conditions, as well as in modulating the secondary immune response by co-stimulating

memory T-cell function (By similarity).[UniProtKB/Swiss-Prot Function]

Protein Families: Transmembrane

Protein Pathways: Cell adhesion molecules (CAMs)

Product images:



Coomassie blue staining of purified ICOSLG protein (Cat# [TP308975]). The protein was produced from HEK293T cells transfected with ICOSLG cDNA clone (Cat# [RC208975]) using MegaTran 2.0 (Cat# [TT210002]).