

# **Product datasheet for TP309182M**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### ZADH1 (PTGR2) (NM\_152444) Human Recombinant Protein

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Recombinant protein of human prostaglandin reductase 2 (PTGR2), 100 μg

Species: Human
Expression Host: HEK293T

**Expression cDNA Clone** >RC209182 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MIVQRVVLNSRPGKNGNPVAENFRMEEVYLPDNINEGQVQVRTLYLSVDPYMRCRMNEDTGTDYITPWQL SQVVDGGGIGIIEESKHTNLTKGDFVTSFYWPWQTKVILDGNSLEKVDPQLVDGHLSYFLGAIGMPGLTS LIGIQEKGHITAGSNKTMVVSGAAGACGSVAGQIGHFLGCSRVVGICGTHEKCILLTSELGFDAAINYKK DNVAEQLRESCPAGVDVYFDNVGGNISDTVISQMNENSHIILCGQISQYNKDVPYPPPLSPAIEAIQKER NITRERFLVLNYKDKFEPGILQLSQWFKEGKLKIKETVINGLENMGAAFQSMMTGGNIGKQIVCISEEIS

L

**TRTRPLEQKLISEEDLAANDILDYKDDDDKV** 

Tag: C-Myc/DDK

Predicted MW: 38.3 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

**Preparation:** Recombinant protein was captured through anti-DDK affinity column followed by conventional

chromatography steps.

**Note:** For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

**Stability:** Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

**RefSeg:** NP 689657

**Locus ID:** 145482



#### ZADH1 (PTGR2) (NM\_152444) Human Recombinant Protein - TP309182M

**UniProt ID:** Q8N8N7, V9HW32

RefSeq Size: 2610

Cytogenetics: 14q24.3 1053 RefSeq ORF:

Synonyms: HEL-S-298; PGR2; ZADH1


**Summary:** This gene encodes an enzyme involved in the metabolism of prostaglandins. The encoded

> protein catalyzes the NADPH-dependent conversion of 15-keto-prostaglandin E2 to 15-keto-13,14-dihydro-prostaglandin E2. This protein may also be involved in regulating activation of the peroxisome proliferator-activated receptor. Alternative splicing results in multiple

transcript variants. [provided by RefSeq, Apr 2009]

**Protein Families:** Druggable Genome

## **Product images:**



Coomassie blue staining of purified PTGR2 protein (Cat# [TP309182]). The protein was produced from HEK293T cells transfected with PTGR2 cDNA clone (Cat# [RC209182]) using

MegaTran 2.0 (Cat# [TT210002]).