

Product datasheet for TP309262

Vitamin D Receptor (VDR) (NM_000376) Human Recombinant Protein

Product data:


Product Type:	Recombinant Proteins
Description:	Recombinant protein of human vitamin D (1,25- dihydroxyvitamin D3) receptor (VDR), transcript variant 1, 20 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC209262 protein sequence Red =Cloning site Green =Tags(s)
	MEAMAASTSLPDPGDFDRNVPRICGVCGRATGFHFNAMTCEGCKGFFRRSMKRKALFTCPFNGDCRIT K DNRRHCQACRLKRCVDIGMMKEFILTDEEVQRKREMILKRKEEALKDSLRLPKLSEEQQRIIALLDAHH KTYDPTYSDFCQFRPPVRVNDGGGSHPSRPNRHTPSFGDSSSSCSHDHCITSSDMMDDSSSFSNLDLSEE DSDDPSVTLELSQLSMLPHLADLVSYISIQKVIGFAKMIPGFRDLTSEDQIVLLKSSAIEVIMLRSNESFT MDDMSWTCGNQDYKYRVSDVTKAGHSLELIEPLIKFQVGLKLNHHEEHVLLMAICIVSPDRPGVQDAA LIEAIQDRLSNTLQTYIRCRHPPPGSHLLYAKMIQKLADLRSLNEEHKQYRCLSFQPECSMKLTPLVLE VFGNEIS
	TRTRPLEQKLISEEDLAANDILDYKDDDDKV
Tag:	C-Myc/DDK
Predicted MW:	48.1 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.



[View online »](#)

Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_000367
Locus ID:	7421
UniProt ID:	P11473
RefSeq Size:	4669
Cytogenetics:	12q13.11
RefSeq ORF:	1281
Synonyms:	NR111; PPP1R163
Summary:	This gene encodes vitamin D3 receptor, which is a member of the nuclear hormone receptor superfamily of ligand-inducible transcription factors. This receptor also functions as a receptor for the secondary bile acid, lithocholic acid. Downstream targets of vitamin D3 receptor are principally involved in mineral metabolism, though this receptor regulates a variety of other metabolic pathways, such as those involved in immune response and cancer. Mutations in this gene are associated with type II vitamin D-resistant rickets. A single nucleotide polymorphism in the initiation codon results in an alternate translation start site three codons downstream. Alternatively spliced transcript variants encoding different isoforms have been described for this gene. A recent study provided evidence for translational readthrough in this gene, and expression of an additional C-terminally extended isoform via the use of an alternative in-frame translation termination codon. [provided by RefSeq, Jun 2018]
Protein Families:	Druggable Genome, Nuclear Hormone Receptor, Transcription Factors

Product images:



Coomassie blue staining of purified VDR protein (Cat# TP309262). The protein was produced from HEK293T cells transfected with VDR cDNA clone (Cat# [RC209262]) using MegaTran 2.0 (Cat# [TT210002]).