

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TP309326

PSMB5 (NM_002797) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins	
Description:	Recombinant protein of human proteasome (prosome, macropain) subunit, beta type, 5 (PSMB5), transcript variant 1, 20 μg	
Species:	Human	
Expression Host:	HEK293T	
Expression cDNA Clone or AA Sequence:	e >RC209326 protein sequence Red=Cloning site Green=Tags(s)	
	MALASVLERPLPVNQRGFFGLGGRADLLDLGPGSLSDGLSLAAPGWGVPEEPGIEMLHGTTTLAFKFRHG VIVAADSRATAGAYIASQTVKKVIEINPYLLGTMAGGAADCSFWERLLARQCRIYELRNKERISVAAASK LLANMVYQYKGMGLSMGTMICGWDKRGPGLYYVDSEGNRISGATFSVGSGSVYAYGVMDRGYSYDLEVE Q AYDLARRAIYQATYRDAYSGGAVNLYHVREDGWIRVSSDNVADLHEKYSGSTP	
	TRTRPLEQKLISEEDLAANDILDYKDDDDKV	
Tag:	C-Myc/DDK	
Predicted MW:	28.3 kDa	
Concentration:	>0.05 µg/µL as determined by microplate BCA method	
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining	
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol	
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.	
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.	
Storage:	Store at -80°C.	
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.	
RefSeq:	<u>NP 002788</u>	
Locus ID:	5693	



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	PSMB5 (NM_002797) Human Recombinant Protein – TP309326
UniProt ID:	<u>P28074</u>
RefSeq Size:	1311
Cytogenetics:	14q11.2
RefSeq ORF:	789
Synonyms:	LMPX; MB1; X
Summary:	The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the proteasome B-type family, also known as the T1B family, that is a 20S core beta subunit in the proteasome. This catalytic subunit is not present in the immunoproteasome and is replaced by catalytic subunit 3i (proteasome beta 8 subunit). Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jan 2009]
Protein Families:	Protease
Protein Pathway	s: Proteasome

Product images:

188	_	
98	-	
62	_	
49	_	
38	_	
28	_	-
17 14 63	Ξ	

Coomassie blue staining of purified PSMB5 protein (Cat# TP309326). The protein was produced from HEK293T cells transfected with PSMB5 cDNA clone (Cat# [RC209326]) using MegaTran 2.0 (Cat# [TT210002]).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US