

Product datasheet for **TP309462L**

ATP6V1B1 (NM_001692) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human ATPase, H ⁺ transporting, lysosomal 56/58kDa, V1 subunit B1 (ATP6V1B1), 1 mg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC209462 protein sequence Red =Cloning site Green =Tags(s)

MAMEIDSRPGGLPGSSCNLGAAREHMQAVTRNYITHPRVYRTVCSVNGPLVLDLDRVKFAQYAEIVHFTL
PDGTQRSGQVLEVAGTKAIVQVFEGTSGIDARKTTCEFTGDILRTPVSEDMLGRVFNGSGKPIDKGPVVM
AEDFLDINGQPINPHSRIYPEEMIQTGISPIDVMNSIARGQKIPIFSAAGLPHNEIAAQICRQAGLVKKS
KAVLDYHDDNFAIVFAAMGVNMETARFFKSDFEQNGTGMNVCLFLNLANPTIERIITPRALTTAEFLA
YQCEKHVLVILDMSSYAEALREVSAAAREEVPGRRGFPGYMYTDLATYERAGRVEGRGGSITQIPILTM
PNDDITHPIPDLTGFITEGQIYVDRQLHNRQIYPPINVLPSLSRLMKSAIGEGMTRKDHGDVSNQLYACY
AIGKDVQAMKAVVGEEALTSEDLLYLEFLQKFEKNFINQGPYENRSVFESLDLGGWKLRLRIFPKEMLKRI
QAVIDEFYSREGALQDLAPDTAL

TRTRPLEQKLISEEDLAANDILDYKDDDDKV


Tag:	C-Myc/DDK
Predicted MW:	56.7 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.



[View online »](#)

Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_001683
Locus ID:	525
UniProt ID:	P15313
RefSeq Size:	1956
Cytogenetics:	2p13.3
RefSeq ORF:	1539
Synonyms:	ATP6B1; DRTA2; RTA1B; VATB; VMA2; VPP3
Summary:	This gene encodes a component of vacuolar ATPase (V-ATPase), a multisubunit enzyme that mediates acidification of eukaryotic intracellular organelles. V-ATPase dependent organelle acidification is necessary for such intracellular processes as protein sorting, zymogen activation, receptor-mediated endocytosis, and synaptic vesicle proton gradient generation. V-ATPase is composed of a cytosolic V1 domain and a transmembrane V0 domain. The V1 domain consists of three A and three B subunits, two G subunits plus the C, D, E, F, and H subunits. The V1 domain contains the ATP catalytic site. The V0 domain consists of five different subunits: a, c, c', c'', and d. Additional isoforms of many of the V1 and V0 subunit proteins are encoded by multiple genes or alternatively spliced transcript variants. This encoded protein is one of two V1 domain B subunit isoforms and is found in the kidney. Mutations in this gene cause distal renal tubular acidosis associated with sensorineural deafness. [provided by RefSeq, Jul 2008]
Protein Families:	Druggable Genome
Protein Pathways:	Epithelial cell signaling in Helicobacter pylori infection, Metabolic pathways, Oxidative phosphorylation, Vibrio cholerae infection

Product images:



Coomassie blue staining of purified ATP6V1B1 protein (Cat# [TP309462]). The protein was produced from HEK293T cells transfected with ATP6V1B1 cDNA clone (Cat# [RC209462]) using MegaTran 2.0 (Cat# [TT210002]).