

Product datasheet for TP309500L

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

THOC7 (NM_025075) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human THO complex 7 homolog (Drosophila) (THOC7), 1 mg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC209500 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MGAVTDDEVIRKRLLIDGDGAGDDRRINLLVKSFIKWCNSGSQEEGYSQYQRMLSTLSQCEFSMGKTLLV YDMNLREMENYEKIYKEIECSIAGAHEKIAECKKQILQAKRIRKNRQEYDALAKVIQHHPDRHETLKELE ALGKELEHLSHIKESVEDKLELRRKQFHVLLSTIHELQQTLENDEKLSEVEEAQEASMETDPKP

TRTRPLEQKLISEEDLAANDILDYKDDDDK**V**

Tag: C-Myc/DDK
Predicted MW: 23.6 kDa

Concentration: >0.05 μg/μL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 079351

 Locus ID:
 80145

 UniProt ID:
 Q619Y2

 RefSeq Size:
 1027



ORÏGENE

Cytogenetics: 3p14.1

RefSeq ORF: 612

Synonyms: fSAP24; hTREX30; NIF3L1BP1


Summary: Required for efficient export of polyadenylated RNA. Acts as component of the THO

> subcomplex of the TREX complex which is thought to couple mRNA transcription, processing and nuclear export, and which specifically associates with spliced mRNA and not with unspliced pre-mRNA. TREX is recruited to spliced mRNAs by a transcription-independent mechanism, binds to mRNA upstream of the exon-junction complex (EJC) and is recruited in a splicing- and cap-dependent manner to a region near the 5' end of the mRNA where it

> functions in mRNA export to the cytoplasm via the TAP/NFX1 pathway.[UniProtKB/Swiss-Prot

Function]

Product images:



Coomassie blue staining of purified THOC7 protein (Cat# [TP309500]). The protein was produced from HEK293T cells transfected with THOC7 cDNA clone (Cat# [RC209500]) using MegaTran 2.0 (Cat# [TT210002]).