

Product datasheet for TP309701

Influenza Virus NS1A Binding Protein (IVNS1ABP) (NM_006469) Human Recombinant Protein

Product data:


Product Type:	Recombinant Proteins
Description:	Recombinant protein of human influenza virus NS1A binding protein (IVNS1ABP), 20 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC209701 protein sequence Red=Cloning site Green=Tags(s)
	<p>MIPNGYLMFEDENFIESSVAKLNALRKSGQFCDVRLQVCGHEMLAHRAVLACCSPYLFEIFNSDSDPHGI SHVKFDDLNPEAVEVLLNYAYTAQLKADKELVKDVVYSAAKKMKMDRVKQVCGDYLLSRMDVTSCISYRNF ASCMGDSRLLNKVDAYIQEHLLQISEEEFLKLPRLKLEVMLEDNVCLPSNGKLYTKVINWVQRSIWENG DSLEELMGEVQTLYYSADHKLLDGNLLDGQAEVFGSDDDDHIQFVQKKPPRENGHKQISSSSTGCLSSPNA TVQSPKHEWKIVASEKTSNNTYLCLAVLDGIFCVIFLHGRNSPQSSPTSTPKLSKLSLFEMQQDELIEKP MSPMQYARSLGTAEMNGKLIAGGYNREECLRTVECYNPHTDHWSFLAPMRTPRARFQMAVLMGQLYV GGSNGHSDDLSCGEMYDSNIDDWIPVELRTRNRCNAGVCALNGKLYVGGSDPYGQKGLKNCDVFDPVTK LWTSCAPLNIRRHQSAVCELGGYLYIIGGAESWNCLNTVERYNPENNTWTLIAPMNVARRGAGVAVLNGK LFVCGGFDGSHAISCVEMYDPTREWKMMGNMTPRSNAGIATVGNTIYAVGGFDGNEFLNTEVYNLES NEWSPYTKIFQF</p> <p>TRTRPLEQKLISEEDLAANDILDYKDDDDKV</p>
Tag:	C-Myc/DDK
Predicted MW:	71.5 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.



[View online »](#)

Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_006460
Locus ID:	10625
UniProt ID:	Q9Y6Y0
RefSeq Size:	4113
Cytogenetics:	1q25.3
RefSeq ORF:	1929
Synonyms:	ARA3; FLARA3; HSPC068; IMD70; KLHL39; ND1; NS-1; NS1-BP; NS1BP
Summary:	Involved in many cell functions, including pre-mRNA splicing, the aryl hydrocarbon receptor (AHR) pathway, F-actin organization and protein ubiquitination. Plays a role in the dynamic organization of the actin skeleton as a stabilizer of actin filaments by association with F-actin through Kelch repeats (By similarity). Protects cells from cell death induced by actin destabilization (By similarity). Functions as modifier of the AHR/Aryl hydrocarbon receptor pathway increasing the concentration of AHR available to activate transcription (PubMed:16582008). In addition, functions as a negative regulator of BCR(KLHL20) E3 ubiquitin ligase complex to prevent ubiquitin-mediated proteolysis of PML and DAPK1, two tumor suppressors (PubMed:25619834). Inhibits pre-mRNA splicing (in vitro) (PubMed:9696811). [UniProtKB/Swiss-Prot Function]
Protein Families:	Transcription Factors

Product images:



Coomassie blue staining of purified IVNS1ABP protein (Cat# TP309701). The protein was produced from HEK293T cells transfected with IVNS1ABP cDNA clone (Cat# [RC209701]) using MegaTran 2.0 (Cat# [TT210002]).