

Product datasheet for TP309810

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

C13orf15 (RGCC) (NM_014059) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human chromosome 13 open reading frame 15 (C13orf15), 20 μg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC209810 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MKQPAAQGSPAAAAAAAAAALDSAAAEDLSDALCEFDAVLADFASPFHERHFHYEEHLERMKRRSSASVSD

SSGFSDSESADSLYRNSFSFSDEKLNSPTDSTPALLSATVTPQKAKLGDTKELEAFIADLDKTLASM

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK
Predicted MW: 14.4 kDa

Concentration: >0.05 μg/μL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 054778

 Locus ID:
 28984

 UniProt ID:
 Q9H4X1

 RefSeq Size:
 1126

 Cytogenetics:
 13q14.11



C13orf15 (RGCC) (NM_014059) Human Recombinant Protein - TP309810

RefSeq ORF: 411


Synonyms: bA157L14.2; C13orf15; RGC-32; RGC32

Summary: This gene is thought to regulate cell cycle progression. It is induced by p53 in response to

DNA damage, or by sublytic levels of complement system proteins that result in activation of the cell cycle. The encoded protein localizes to the cytoplasm during interphase and to centrosomes during mitosis. The protein forms a complex with polo-like kinase 1. The protein also translocates to the nucleus in response to treatment with complement system proteins, and can associate with and increase the kinase activity of cell division cycle 2 protein. In different assays and cell types, overexpression of this protein has been shown to activate or

suppress cell cycle progression. [provided by RefSeq, Jul 2008]

Product images:



Coomassie blue staining of purified RGCC protein (Cat# TP309810). The protein was produced from HEK293T cells transfected with RGCC cDNA clone (Cat# [RC209810]) using MegaTran 2.0 (Cat# [TT210002]).