

Product datasheet for TP309976L

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

CN: techsupport@origene.cn

Proteasome subunit alpha type 6 (PSMA6) (NM_002791) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human proteasome (prosome, macropain) subunit, alpha type, 6

(PSMA6), 1 mg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC209976 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MSRGSSAGFDRHITIFSPEGRLYQVEYAFKAINQGGLTSVAVRGKDCAVIVTQKKVPDKLLDSSTVTHLF KITENIGCVMTGMTADSRSQVQRARYEAANWKYKYGYEIPVDMLCKRIADISQVYTQNAEMRPLGCCMIL IGIDEEQGPQVYKCDPAGYYCGFKATAAGVKQTESTSFLEKKVKKKFDWTFEQTVETAITCLSTVLSIDF

KPSEIEVGVVTVENPKFRILTEAEIDAHLVALAERD

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Predicted MW: 27.2 kDa

Concentration: $>0.05 \mu g/\mu L$ as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 002782

Locus ID: 5687





Proteasome subunit alpha type 6 (PSMA6) (NM_002791) Human Recombinant Protein – TP309976L

UniProt ID: <u>P60900</u>, <u>A0A140VK44</u>

RefSeq Size: 1091 Cytogenetics: 14q13.2 RefSeq ORF: 738

Synonyms: IOTA; p27K; PROS27

Summary: The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S


core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the peptidase T1A family, that is a 20S core alpha subunit. Multiple transcript variants encoding several different isoforms have been found for this gene.

A pseudogene has been identified on the Y chromosome. [provided by RefSeq, Aug 2013]

Protein Families: Druggable Genome, Protease, Stem cell - Pluripotency

Protein Pathways: Proteasome

Product images:



Coomassie blue staining of purified PSMA6 protein (Cat# [TP309976]). The protein was produced from HEK293T cells transfected with PSMA6 cDNA clone (Cat# [RC209976]) using MegaTran 2.0 (Cat# [TT210002]).