

# **Product datasheet for TP310027M**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### PYCR1 (NM\_006907) Human Recombinant Protein

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Recombinant protein of human pyrroline-5-carboxylate reductase 1 (PYCR1), transcript variant

1, 100 μg

Species: Human
Expression Host: HEK293T

**Expression cDNA Clone** >RC210027 representing NM\_006907

or AA Sequence: Red=Cloning site Green=Tags(s)

MSVGFIGAGQLAFALAKGFTAAGVLAAHKIMASSPDMDLATVSALRKMGVKLTPHNKETVQHSDVLFLAV

KPHIIPFILDEIGADIEDRHIVVSCAAGVTISSIEKKLSAFRPAPRVIRCMTNTPVVVREGATVYATGTH

AQVEDGRLMEQLLSSVGFCTEVEEDLIDAVTGLSGSGPAYAFTALDALADGGVKMGLPRRLAVRLGAQAL LGAAKMLLHSEQHPGQLKDNVSSPGGATIHALHVLESGGFRSLLINAVEASCIRTRELQSMADQEQVSPA

AIKKTILDKVKLDSPAGTALSPSGHTKLLPRSLAPAGKD

**TRTRPL**EQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Predicted MW: 33.2 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

**Preparation:** Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

**Note:** For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

**Stability:** Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 008838

Locus ID: 5831



# PYCR1 (NM\_006907) Human Recombinant Protein - TP310027M

UniProt ID: <u>P32322</u>, <u>A0A024R8U9</u>, <u>Q8TBX0</u>

RefSeq Size: 2059

Cytogenetics: 17q25.3

RefSeq ORF: 957

Synonyms: ARCL2B; ARCL3B; P5C; P5CR; PIG45; PP222; PRO3; PYCR

Summary: This gene encodes an enzyme that catalyzes the NAD(P)H-dependent conversion of pyrroline-

5-carboxylate to proline. This enzyme may also play a physiologic role in the generation of


NADP(+) in some cell types. The protein forms a homopolymer and localizes to the

mitochondrion. Alternative splicing results in multiple transcript variants. [provided by RefSeq,

Aug 2013]

**Protein Pathways:** Arginine and proline metabolism, Metabolic pathways

# **Product images:**



Coomassie blue staining of purified PYCR1 protein (Cat# [TP310027]). The protein was produced from HEK293T cells transfected with PYCR1 cDNA clone (Cat# [RC210027]) using

MegaTran 2.0 (Cat# [TT210002]).