

Product datasheet for **TP310035M**

PCBP2 (NM_005016) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human poly(rC) binding protein 2 (PCBP2), transcript variant 1, 100 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC210035 protein sequence Red =Cloning site Green =Tags(s)

MDTGVIIEGGLNVTLRLLMHGKEVGSIIIGKKGESVKKMREESGARINISEGNCPERIITLAGPTNAIFK
AFAMIIDKLEEDISSSMTNSTAASRPPVTLRLVVPASQCGSLIGKGGCKIKEIRESTGAQVQVAGDMLPN
STERAITIAGIPQSIIIECVKQICVVMLETLSQSPPKGV TIPYRKPSSSPVIFAGGQDRYSTGSDSASFP
HTTSPMCLNPDLEGPPLEAYTIQGQYAIQPDLTCLKLHQLAMQQSHFPMTHGNTGFSGIESSSPEVKGYWA
GLDASAQTTSHELTIPNDLIGCIIGRQGAKINEIRQMSGAQIKIANPVEGSTDRQVTITGSAASISLAQY
LINVRLSSETGGMGSS

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag:	C-Myc/DDK
Predicted MW:	38.5 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Bioactivity:	Pull-down assay (PMID: 25855805)
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_005007



[View online »](#)

Locus ID: 5094
UniProt ID: [Q15366](#), [A0A384N6B9](#)
RefSeq Size: 3187
Cytogenetics: 12q13.13
RefSeq ORF: 1098
Synonyms: hnRNP-E2; HNRNPE2; HNRPE2


Summary: The protein encoded by this gene appears to be multifunctional. Along with PCBP-1 and hnRNPK, it is one of the major cellular poly(rC)-binding proteins. The encoded protein contains three K-homologous (KH) domains which may be involved in RNA binding. Together with PCBP-1, this protein also functions as a translational coactivator of poliovirus RNA via a sequence-specific interaction with stem-loop IV of the IRES, promoting poliovirus RNA replication by binding to its 5'-terminal cloverleaf structure. It has also been implicated in translational control of the 15-lipoxygenase mRNA, human papillomavirus type 16 L2 mRNA, and hepatitis A virus RNA. The encoded protein is also suggested to play a part in formation of a sequence-specific alpha-globin mRNP complex which is associated with alpha-globin mRNA stability. This multiexon structural mRNA is thought to be retrotransposed to generate PCBP-1, an intronless gene with functions similar to that of PCBP2. This gene and PCBP-1 have paralogous genes (PCBP3 and PCBP4) which are thought to have arisen as a result of duplication events of entire genes. This gene also has two processed pseudogenes (PCBP2P1 and PCBP2P2). Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jan 2018]

Product images:

HS21	-	+	+	+	+	-
beads	-	+	+	+	+	+
PCBP2	+	+	+	+	+	+
TMPyP4	-	0				20x



The i-motif stabilizing ligand (TMPyP4) displaces telomere-associated human poly-C binding protein (PCBP) from human C-rich telomeric DNA. Biotinylated C-rich telomeric DNA HS21 oligo was bound to streptavidin magnetic beads and incubated with PCBP2 (OriGene [TP310035]) in the absence or presence of TMPyP4 at different concentrations. The bound PCBP2 was eluted and analyzed in Western blot with an anti-PCBP2 antibody. Figure cited from Nucleic Acids Res, PMID: 25855805



Coomassie blue staining of purified PCBP2 protein (Cat# [TP310035]). The protein was produced from HEK293T cells transfected with PCBP2 cDNA clone (Cat# [RC210035]) using MegaTran 2.0 (Cat# [TT210002]).