

Product datasheet for **TP310190**

UBASH3A (NM_018961) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human ubiquitin associated and SH3 domain containing, A (UBASH3A), transcript variant 1, 20 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC210190 representing NM_018961 Red =Cloning site Green =Tags(s)

MAAGETQLYAKVSNKLGKRSSPSLLEPLLAMGFPVHTALKALAATGRKTAEEALAWLHDHCNDPSLDDPI
PQEYALFLCPTGPLELQEFWRESKRQCAKNRAHEVFPHTLDCFFTCEDQKVECLYEALKRAGDRLLG
SFPTAVPLALHSSISYLGFFVSGSPADVIREFAMTFATEASLLAGTSVSRWFIFSQVPGHGPNLRLSNLT
RASFSVSHYILQKYCSVKPCTKQLHLTLAHKFYPHHQRTLEQLARAIPLGHSCQWTAALYSRDMRFVHYQT
LRVLFQYKPNVDELTLSPGDYIFVDPTQQDEASEGWVIGISQRTGCRGFLPENYTDRASESDTWVKHRM
YTFSLATDLNSRKDGEASSRCSGEFLPQTARSLSSLQALQATVARKSVLVVRHGERVDQIFGKAWLQQCS
TPDGKYRDLNFP CSLPRRSRGIKDFENDPPLSSCGIFQSR IAGDALLDSGIRISSVFASPALRCVQTA
KLILEELKLEKKIKIRVEPGIFEWTKWEAGKTTPTLMSLEELKEANFNIDTDYRPAFPLSALMPAESYQE
YMDRCTASMVQIVNTCPQDTGVILIVSHGSTLDSCTRPLLGLPPRECGDFAQLVRKIPSLGMCFCEENKE
EGKWELVNPVKTLTHGANAAFNRNWNISGN

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag:	C-Myc/DDK
Predicted MW:	73.9 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.



[View online >](#)

Stability: Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: [NP_061834](#)

Locus ID: 53347

UniProt ID: [P57075](#)

RefSeq Size: 2492

Cytogenetics: 21q22.3


RefSeq ORF: 1983

Synonyms: CLIP4; STS-2; TULA; TULA-1

Summary: This gene encodes one of two family members belonging to the T-cell ubiquitin ligand (TULA) family. Both family members can negatively regulate T-cell signaling. This family member can facilitate growth factor withdrawal-induced apoptosis in T cells, which may occur via its interaction with AIF, an apoptosis-inducing factor. Alternative splicing of this gene results in multiple transcript variants. [provided by RefSeq, Aug 2011]

Protein Families: Druggable Genome

Product images:



Coomassie blue staining of purified UBASH3A protein (Cat# TP310190). The protein was produced from HEK293T cells transfected with UBASH3A cDNA clone (Cat# [RC210190]) using MegaTran 2.0 (Cat# [TT210002]).