

Product datasheet for TP310311L

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

CN: techsupport@origene.cn

Retinoid X Receptor alpha (RXRA) (NM_002957) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human retinoid X receptor, alpha (RXRA), 1 mg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC210311 representing NM_002957 **or AA Sequence:** Red=Cloning site Green=Tags(s)

MDTKHFLPLDFSTQVNSSLTSPTGRGSMAAPSLHPSLGPGIGSPGQLHSPISTLSSPINGMGPPFSVISS PMGPHSMSVPTTPTLGFSTGSPQLSSPMNPVSSSEDIKPPLGLNGVLKVPAHPSGNMASFTKHICAICGD RSSGKHYGVYSCEGCKGFFKRTVRKDLTYTCRDNKDCLIDKRQRNRCQYCRYQKCLAMGMKREAVQEERQ RGKDRNENEVESTSSANEDMPVERILEAELAVEPKTETYVEANMGLNPSSPNDPVTNICQAADKQLFTLV EWAKRIPHFSELPLDDQVILLRAGWNELLIASFSHRSIAVKDGILLATGLHVHRNSAHSAGVGAIFDRVL TELVSKMRDMQMDKTELGCLRAIVLFNPDSKGLSNPAEVEALREKVYASLEAYCKHKYPEQPGRFAKLLL

RLPALRSIGLKCLEHLFFFKLIGDTPIDTFLMEMLEAPHQMT

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK
Predicted MW: 50.6 kDa

Concentration: >0.05 μg/μL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by conventional

chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 002948





Locus ID: 6256

UniProt ID: P19793, F1D8Q5, Q6P3U7

RefSeg Size: 5449 Cytogenetics: 9q34.2 RefSeq ORF: 1386 Synonyms: NR2B1

Summary: Retinoid X receptors (RXRs) and retinoic acid receptors (RARs) are nuclear receptors that

> mediate the biological effects of retinoids by their involvement in retinoic acid-mediated gene activation. These receptors function as transcription factors by binding as homodimers or heterodimers to specific sequences in the promoters of target genes. The protein encoded by

this gene is a member of the steroid and thyroid hormone receptor superfamily of transcriptional regulators. Alternative splicing of this gene results in multiple transcript


variants. [provided by RefSeq, May 2014]

Protein Families: Druggable Genome, Nuclear Hormone Receptor, Transcription Factors

Protein Pathways: Adipocytokine signaling pathway, Non-small cell lung cancer, Pathways in cancer, PPAR

signaling pathway, Small cell lung cancer, Thyroid cancer

Product images:



Coomassie blue staining of purified RXRA protein (Cat# [TP310311]). The protein was produced from HEK293T cells transfected with RXRA cDNA clone (Cat# [RC210311]) using MegaTran 2.0

(Cat# [TT210002]).