

# **Product datasheet for TP310316**

# OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Cyclin D2 (CCND2) (NM\_001759) Human Recombinant Protein

#### **Product data:**

**Product Type:** Recombinant Proteins

**Description:** Recombinant protein of human cyclin D2 (CCND2), 20 μg

Species: Human
Expression Host: HEK293T

**Expression cDNA Clone** >RC210316 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MELLCHEVDPVRRAVRDRNLLRDDRVLQNLLTIEERYLPQCSYFKCVQKDIQPYMRRMVATWMLEVCEEQ KCEEEVFPLAMNYLDRFLAGVPTPKSHLQLLGAVCMFLASKLKETSPLTAEKLCIYTDNSIKPQELLEWE LVVLGKLKWNLAAVTPHDFIEHILRKLPQQREKLSLIRKHAQTFIALCATDFKFAMYPPSMIATGSVGAA ICGLQQDEEVSSLTCDALTELLAKITNTDVDCLKACQEQIEAVLLNSLQQYRQDQRDGSKSEDELDQAST

PTDVRDIDL

**TRTRPL**EQKLISEEDLAANDILDYKDDDDK**V** 

Tag: C-Myc/DDK

Predicted MW: 32.9 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

**Preparation:** Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

**Note:** For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

**RefSeq:** NP 001750

Locus ID: 894



### Cyclin D2 (CCND2) (NM\_001759) Human Recombinant Protein - TP310316

**UniProt ID:** P30279 RefSeq Size: 6531

**Cytogenetics:** 12p13.32

RefSeq ORF: 867

KIAK0002; MPPH3 Synonyms:

**Summary:** The protein encoded by this gene belongs to the highly conserved cyclin family, whose

members are characterized by a dramatic periodicity in protein abundance through the cell

cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct

expression and degradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a complex with CDK4 or CDK6 and functions as a regulatory subunit of the complex, whose activity is required for cell cycle G1/S transition. This protein has been shown to interact with and be involved in the phosphorylation of tumor suppressor protein Rb. Knockout studies of the homologous gene in mouse suggest the essential roles of this gene in ovarian granulosa and germ cell proliferation. High level expression of this gene was observed in ovarian and testicular tumors. Mutations in this gene are associated with megalencephaly-polymicrogyria-polydactyly-hydrocephalus syndrome 3 (MPPH3). [provided


by RefSeq, Sep 2014]

**Protein Families:** Druggable Genome

**Protein Pathways:** Cell cycle, Focal adhesion, Jak-STAT signaling pathway, p53 signaling pathway, Wnt signaling

pathway

## **Product images:**



Coomassie blue staining of purified CCND2 protein (Cat# TP310316). The protein was produced from HEK293T cells transfected with CCND2 cDNA clone (Cat# [RC210316]) using

MegaTran 2.0 (Cat# [TT210002]).