

Product datasheet for TP310404L

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Proteinase Activated Receptor 4 (F2RL3) (NM 003950) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human coagulation factor II (thrombin) receptor-like 3 (F2RL3), 1 mg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC210404 representing NM_003950 or AA Sequence: Red=Cloning site Green=Tags(s)

MWGRLLLWPLVLGFSLSGGTQTPSVYDESGSTGGGDDSTPSILPAPRGYPGQVCANDSDTLELPDSSRAL LLGWVPTRLVPALYGLVLVVGLPANGLALWVLATQAPRLPSTMLLMNLAAADLLLALALPPRIAYHLRGQ RWPFGEAACRLATAALYGHMYGSVLLLAAVSLDRYLALVHPLRARALRGRRLALGLCMAAWLMAAALALP LTLQRQTFRLARSDRVLCHDALPLDAQASHWQPAFTCLALLGCFLPLLAMLLCYGATLHTLAASGRRYGH ALRLTAVVLASAVAFFVPSNLLLLLHYSDPSPSAWGNLYGAYVPSLALSTLNSCVDPFIYYYVSAEFRDK

VRAGLFQRSPGDTVASKASAEGGSRGMGTHSSLLQ

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Predicted MW: 36.2 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 003941

Locus ID: 9002





UniProt ID: Q96RIO

RefSeq Size: 2731

Cytogenetics: 19p13.11
RefSeq ORF: 1155
Synonyms: PAR4

Summary: This gene encodes a member of the protease-activated receptor subfamily, part of the G-


protein coupled receptor 1 family of proteins. The encoded receptor is proteolytically processed to reveal an extracellular N-terminal tethered ligand that binds to and activates the receptor. This receptor plays a role in blood coagulation, inflammation and response to pain.

Hypomethylation at this gene may be associated with lung cancer in human patients.

[provided by RefSeq, Sep 2016]

Protein Families: Druggable Genome, GPCR, Transmembrane
Protein Pathways: Neuroactive ligand-receptor interaction

Product images:



Coomassie blue staining of purified F2RL3 protein (Cat# [TP310404]). The protein was produced from HEK293T cells transfected with F2RL3 cDNA clone (Cat# [RC210404]) using MegaTran 2.0 (Cat# [TT210002]).