

Product datasheet for TP310452M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Proteasome subunit beta type 2 (PSMB2) (NM_002794) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human proteasome (prosome, macropain) subunit, beta type, 2

(PSMB2), 100 µg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC210452 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MEYLIGIQGPDYVLVASDRVAASNIVQMKDDHDKMFKMSEKILLLCVGEAGDTVQFAEYIQKNVQLYKMR NGYELSPTAAANFTRRNLADCLRSRTPYHVNLLLAGYDEHEGPALYYMDYLAALAKAPFAAHGYGAFLTL

SILDRYYTPTISRERAVELLRKCLEELQKRFILNLPTFSVRIIDKNGIHDLDNISFPKQGS

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Predicted MW: 22.7 kDa

Concentration: $>0.05 \mu g/\mu L$ as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 002785

Locus ID: 5690

UniProt ID: P49721, A0A140V|S6





Proteasome subunit beta type 2 (PSMB2) (NM_002794) Human Recombinant Protein – TP310452M

RefSeq Size: 4759

Cytogenetics: 1p34.3
RefSeq ORF: 603
Synonyms: HC7-I

Summary: The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S


core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the proteasome B-type family, also known as the T1B family, that is a 20S core beta subunit. Multiple alternatively spliced transcript variants encoding distinct

isoforms have been found for this gene. [provided by RefSeq, Dec 2010]

Protein Families: Druggable Genome, Protease

Protein Pathways: Proteasome

Product images:



Coomassie blue staining of purified PSMB2 protein (Cat# [TP310452]). The protein was produced from HEK293T cells transfected with PSMB2 cDNA clone (Cat# [RC210452]) using MegaTran 2.0 (Cat# [TT210002]).