

Product datasheet for TP310576

MACROH2A1 (NM_004893) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human H2A histone family, member Y (H2AFY), transcript variant 2, 20 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC210576 protein sequence Red=Cloning site Green=Tags(s)

MSSRGGKKKSTKTSRSAKAGVIFPVGRMLRYIKKGHPKYRIGVVGAPVYMAAVLEYLTAEILELAGNAARD
NKKGRVTPRHILLAVANDEELNQLLKGVTIASGGVLPNIHPELLAKKRGSKGKLEAITPPPAKAKSPS
QKKPVSKKAGGKKGARKSKKKQGEVSKAASADSTTEGTPADGFTVLSTKSLFLGQKLNLIHSEISNLGAF
EVEAIINPTNADIDLKDDLGNLTLEKGGKEFVEAVLELRKKNNGPLEVAGAAVSAGHGLPAKFVIHCNSPV
WGADKCEELLGKTVKNCLALADDKCLKSIAFPSIGSGRNGFPPKQTAAQLILKAISSYFVSTMSSSIKTVY
FVLFDESEIGIYVQEMAKLDAN

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag:	C-Myc/DDK
Predicted MW:	39.3 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_004884




[View online »](#)

Locus ID: 9555
UniProt ID: [O75367](#)
RefSeq Size: 1923
Cytogenetics: 5q31.1
RefSeq ORF: 1116
Synonyms: H2A.y; H2A/y; H2AF12M; H2AFY; MACROH2A1.1; macroH2A1.2; mH2A1

Summary: Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene encodes a replication-independent histone that is a member of the histone H2A family. It replaces conventional H2A histones in a subset of nucleosomes where it represses transcription and participates in stable X chromosome inactivation. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Oct 2015]

Protein Pathways: Systemic lupus erythematosus

Product images:



Coomassie blue staining of purified H2AFY protein (Cat# TP310576). The protein was produced from HEK293T cells transfected with H2AFY cDNA clone (Cat# [RC210576]) using MegaTran 2.0 (Cat# [TT210002]).