

Product datasheet for TP310626M

HSPC142 (BABAM1) (NM_001033549) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human chromosome 19 open reading frame 62 (C19orf62), transcript variant 1, 100 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC210626 protein sequence Red =Cloning site Green =Tags(s)

MEVAEPSSPTEEEEEEHSAEPRPRTRSNPEGAEDRAVGAQASVGSRSEGEGEAASADDGSLNTSGAGP
KSWQVPPPAPEVQIRTPRVNCPKVIICLDLSEEMSLPKLEFNGSKTNALNVSQKMIEMFVRTKHKIDK
SHEFALVVNDDTAWLSGLTSDPRELCSCLYDLETASCSTFNLEGLFSLIQQKTELPTENVQTIPPPYV
VRTILVYSRPPCQPQFSLTEPMKKMFQCPYFFFDVVIHNGTTEEKEEEMSWKDMFAFMGSLDTKGTSYKY
EVALAGPALELHNCMAKLLAHPLQRPCQSHASYSLLLEEDEAIEVEATV

TRTRPLEQKLISEEDLAANDILDYKDDDDKV


Tag:	C-Myc/DDK
Predicted MW:	36.4 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	<u>NP_001028721</u>
Locus ID:	29086



[View online »](#)

UniProt ID:	Q9NWW8
RefSeq Size:	1505
Cytogenetics:	19p13.11
RefSeq ORF:	987
Synonyms:	C19orf62; HSPC142; MERIT40; NBA1
Summary:	<p>Component of the BRCA1-A complex, a complex that specifically recognizes 'Lys-63'-linked ubiquitinated histones H2A and H2AX at DNA lesions sites, leading to target the BRCA1-BARD1 heterodimer to sites of DNA damage at double-strand breaks (DSBs). The BRCA1-A complex also possesses deubiquitinase activity that specifically removes 'Lys-63'-linked ubiquitin on histones H2A and H2AX. In the BRCA1-A complex, it is required for the complex integrity and its localization at DSBs. Component of the BRISC complex, a multiprotein complex that specifically cleaves 'Lys-63'-linked ubiquitin in various substrates (PubMed:24075985, PubMed:26195665). In these 2 complexes, it is probably required to maintain the stability of BABAM2 and help the 'Lys-63'-linked deubiquitinase activity mediated by BRCC3/BRCC36 component. The BRISC complex is required for normal mitotic spindle assembly and microtubule attachment to kinetochores via its role in deubiquitinating NUMA1 (PubMed:26195665). Plays a role in interferon signaling via its role in the deubiquitination of the interferon receptor IFNAR1; deubiquitination increases IFNAR1 activity by enhancing its stability and cell surface expression (PubMed:24075985). Down-regulates the response to bacterial lipopolysaccharide (LPS) via its role in IFNAR1 deubiquitination (PubMed:24075985).[UniProtKB/Swiss-Prot Function]</p>

Product images:



Coomassie blue staining of purified BABAM1 protein (Cat# [TP310626]). The protein was produced from HEK293T cells transfected with BABAM1 cDNA clone (Cat# [RC210626]) using MegaTran 2.0 (Cat# [TT210002]).