

## **Product datasheet for TP310717**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Proteasome 20S alpha 5 (PSMA5) (NM 002790) Human Recombinant Protein

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Recombinant protein of human proteasome (prosome, macropain) subunit, alpha type, 5

(PSMA5), 20 µg

Species: Human
Expression Host: HEK293T

**Expression cDNA Clone** >RC210717 representing NM\_002790

or AA Sequence: Red=Cloning site Green=Tags(s)

MFLTRSEYDRGVNTFSPEGRLFQVEYAIEAIKLGSTAIGIQTSEGVCLAVEKRITSPLMEPSSIEKIVEI

DAHIGCAMSGLIADAKTLIDKARVETQNHWFTYNETMTVESVTQAVSNLALQFGEEDADPGAMSRPFGVA LLFGGVDEKGPQLFHMDPSGTFVQCDARAIGSASEGAQSSLQEVYHKSMTLKEAIKSSLIILKQVMEEKL

NATNIELATVQPGQNFHMFTKEELEEVIKDI

**TRTRPL**EQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

**Predicted MW:** 26.2 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

**Preparation:** Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

**Note:** For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

**RefSeq:** NP 002781

Locus ID: 5686





UniProt ID: <u>P28066</u>, <u>A0A109NGN6</u>

RefSeq Size: 1023 Cytogenetics: 1p13.3 RefSeq ORF: 723

Synonyms: PSC5; ZETA

Summary: The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S


core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the peptidase T1A family, that is a 20S core alpha subunit. Multiple alternatively spliced transcript variants encoding two distinct isoforms have been

found for this gene. [provided by RefSeq, Dec 2010]

**Protein Families:** Druggable Genome, Protease

**Protein Pathways:** Proteasome

## **Product images:**



Coomassie blue staining of purified PSMA5 protein (Cat# TP310717). The protein was produced from HEK293T cells transfected with PSMA5 cDNA clone (Cat# [RC210717]) using MegaTran 2.0 (Cat# [TT210002]).