

Product datasheet for **TP310979L**

PUS10 (NM_144709) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human pseudouridylate synthase 10 (PUS10), 1 mg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC210979 protein sequence Red =Cloning site Green =Tags(s)

MFPLTEENKHVAQLLLNTGTCPRCIFRCGVDFHAPYKLPYKELLNELQKFLETEKDELILEVMNPPPKK
IRLQELEDSIDNLSQNGEGRISVSHVGSTASKNSNLNVCNVCLGILQEFCEKDFIKKVCQKVEASGFET
SLVFSVSFPPQLSVREHAALLVKQEMGKQSLSLGRDDIVQLKEAYKWITHPLFSEELGVPIDGKSLFEV
SVVFAHPETVEDCHFLAACPDCFKPAKNKQSVFTRMAVMKALNKIKEEDFLKQFPCPPNSPKAVCAVLE
IECAHGAVFVAGRYNKYSRNLPQTPWIIDGERKLESSVEELISDHLLAVFKAESFNFSSSGREDVDVRTL
GNRPFPAIELVNPVRHVHTSQEIKELQQKINNSSNKIQVRDLQLVTREIGHMKEGEEEEKTKTYSALIWT
NKAIKKKDIEFLNDIKDLKIDKQKTLPLRVLHRRPLAVRARVIHFMETQYVDEHHFRLHLKTQAGTYIKEFV
HGDFGRTPKNIGSLMNVADILELDVESVDVDWPPALDD

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag:	C-Myc/DDK
Predicted MW:	60.1 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.



[View online »](#)

RefSeq: [NP_653310](#)

Locus ID: 150962

UniProt ID: [Q3MIT2](#)

RefSeq Size: 3820


Cytogenetics: 2p16.1-p15

RefSeq ORF: 1587

Synonyms: CCDC139; DOBI; Hup10

Summary: Pseudouridination, the isomerization of uridine to pseudouridine, is the most common posttranscriptional nucleotide modification found in RNA and is essential for biologic functions such as spliceosome biogenesis. Pseudouridylate synthases, such as PUS10, catalyze pseudouridination of structural RNAs, including transfer, ribosomal, and splicing RNAs. These enzymes also act as RNA chaperones, facilitating the correct folding and assembly of tRNAs (McCleverty et al., 2007 [PubMed 17900615]).[supplied by OMIM, May 2009]

Product images:



Coomassie blue staining of purified PUS10 protein (Cat# [TP310979]). The protein was produced from HEK293T cells transfected with PUS10 cDNA clone (Cat# [RC210979]) using MegaTran 2.0 (Cat# [TT210002]).