

Product datasheet for TP311249L

Acetylcholinesterase (ACHE) (NM_015831) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human acetylcholinesterase (Yt blood group) (ACHE), transcript variant E4-E5, 1 mg
Species:	Human
Expression Host:	HEK293T
Expression cDNA	>RC211249 representing NM_015831
Clone or AA Sequence:	Red=Cloning site Green=Tags(s)

MRPPQCLLHTPSLASPLLLLLLWLLGGGVGAEGREDAELLVTVRGGRLRGIRLKTGGPVSAFLGIPFAE
PPMGPRRFLPPEPKQPWSGVVDATTFQSVCYQYVDTLYPGFEGTEMWNPNRESEDCLYLNVWTPYPRPT
SPTPVLWVIYGGGFYSGASSLDVYDGRFLVQAERTVLVSMNYRVGAFGLALPGSREAPGNVGLLDQRLA
LQWVQENVAAFGGDPTSVTLFGESAGAASVGMHLLSPPSRGLFHRAVLQSGAPNGPWATVGMGEARRRAT
QLAHLVGCPPGGTGGNDELVACLRTRPAQVLVNHEWHVLPQESVFRFSFVPVVDGDFLSDTPEALINAG
DFHGLQVLVGVVKDEGSYFLVYGAPGFSKDNESLISRAEFLAGVRVGPVQVSDLAAEAVLHYTDWLHPE
DPARLREALSDVVDHNVCPVAQLAGRLAAQGARVYAYVFEHRASTLSWPLWMGVPHGYEIEFIFGIPL
DPSRNYTAEKIFAQRLMRYWANFARTGDPNEPRDPKAPQWPPYTAGAQQYVSLDLRPLEVRRGLRAQAC
AFWNRFLPKLLSATASEAPSTCPGFTHGEAAPRPGLPLPLLLLHQLLLLFLSHLRRL

TRTRPLEQKLISEEDLAANDILDYKDDDDKV


Tag:	C-Myc/DDK
Predicted MW:	60.8 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.



[View online »](#)

Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_056646
Locus ID:	43
UniProt ID:	P22303
RefSeq Size:	2978
Cytogenetics:	7q22.1
RefSeq ORF:	1851
Synonyms:	ACEE; ARACHE; N-ACHE; YT
Summary:	Acetylcholinesterase hydrolyzes the neurotransmitter, acetylcholine at neuromuscular junctions and brain cholinergic synapses, and thus terminates signal transmission. It is also found on the red blood cell membranes, where it constitutes the Yt blood group antigen. Acetylcholinesterase exists in multiple molecular forms which possess similar catalytic properties, but differ in their oligomeric assembly and mode of cell attachment to the cell surface. It is encoded by the single ACHE gene, and the structural diversity in the gene products arises from alternative mRNA splicing, and post-translational associations of catalytic and structural subunits. The major form of acetylcholinesterase found in brain, muscle and other tissues is the hydrophilic species, which forms disulfide-linked oligomers with collagenous, or lipid-containing structural subunits. The other, alternatively spliced form, expressed primarily in the erythroid tissues, differs at the C-terminal end, and contains a cleavable hydrophobic peptide with a GPI-anchor site. It associates with the membranes through the phosphoinositide (PI) moieties added post-translationally. AChE activity may constitute a sensitive biomarker of RBC ageing in vivo, and thus, may be of aid in understanding the effects of transfusion[provided by RefSeq, Sep 2019]
Protein Families:	Druggable Genome
Protein Pathways:	Glycerophospholipid metabolism

Product images:



Coomassie blue staining of purified ACHE protein (Cat# [TP311249]). The protein was produced from HEK293T cells transfected with ACHE cDNA clone (Cat# [RC211249]) using MegaTran 2.0 (Cat# [TT210002]).