

## **Product datasheet for TP311268M**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## NACA (NM\_005594) Human Recombinant Protein

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Recombinant protein of human nascent polypeptide-associated complex alpha subunit

(NACA), transcript variant 3, 100 μg

Species: Human
Expression Host: HEK293T

**Expression cDNA Clone** >RC211268 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MPGEATETVPATEQELPQPQAETGSGTESDSDESVPELEEQDSTQATTQQAQLAAAAEIDEEPVSKAKQS RSEKKARKAMSKLGLRQVTGVTRVTIRKSKNILFVITKPDVYKSPASDTYIVFGEAKIEDLSQQAQLAAA EKFKVQGEAVSNIQENTQTPTVQEESEEEEVDETGVEVKDIELVMSQANVSRAKAVRALKNNSNDIVNAI

MELTM

**TRTRPLEQKLISEEDLAANDILDYKDDDDKV** 

Tag: C-Myc/DDK
Predicted MW: 23.2 kDa

**Concentration:** >0.05 μg/μL as determined by microplate BCA method

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

**Preparation:** Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

**Note:** For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

**RefSeq:** NP 005585

**Locus ID:** 4666





UniProt ID: Q13765
RefSeq Size: 1112
Cytogenetics: 12q13.3
RefSeq ORF: 645


Synonyms: HSD48; NAC-alpha; NACA1; skNAC

Summary: This gene encodes a protein that associates with basic transcription factor 3 (BTF3) to form

the nascent polypeptide-associated complex (NAC). This complex binds to nascent proteins that lack a signal peptide motif as they emerge from the ribosome, blocking interaction with the signal recognition particle (SRP) and preventing mistranslocation to the endoplasmic reticulum. This protein is an IgE autoantigen in atopic dermatitis patients. Alternative splicing results in multiple transcript variants, but the full length nature of some of these variants, including those encoding very large proteins, has not been determined. There are multiple pseudogenes of this gene on different chromosomes. [provided by RefSeq, Feb 2016]

**Protein Families:** Druggable Genome, Transcription Factors

## **Product images:**



Coomassie blue staining of purified NACA protein (Cat# [TP311268]). The protein was produced from HEK293T cells transfected with NACA cDNA clone (Cat# [RC211268]) using MegaTran 2.0 (Cat# [TT210002]).