

Product datasheet for TP311377M

Cytochrome p450 2C19 (CYP2C19) (NM_000769) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human cytochrome P450, family 2, subfamily C, polypeptide 19 (CYP2C19), 100 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC211377 representing NM_000769 Red=Cloning site Green=Tags(s)
	<p>MDPFVVLVLCLSCLLLLSIWRQSSGRGKLP PGPTPLPVIGNILQIDIKDVSLSLTNLSKIYGPVFTLYFG LERMVVLHGYEVVKEALIDLGEFEFSGRGHFPLAERANRGFGIVFSNGKRWKEIRRFSLMTLRNFGMGKRS IEDRVQEEARCLVEELRKT KASPCDPTFILGCAPCNVICSIIFQKRFDYKDQQLNLMEKLNENIRIVST PWIQCNNFPTIIDYFPGTHNKLLKNLAFMESDILEKVKEHQESMDINNPRDFIDCFLIKMEKEKQNQQS EFTIENLVITAADLLGAGTETTSTLRYALLLLKHPEVTAKVQEEIERVGRNRSPCMQDRGHMPYTD VHEVQRYIDLIPSTLPHAVTCDVKFRNYLIPKGTILTSLTSVLHDNKEFPNPEMFDRHFLDEGGNFK KSNYFMPFSAGKRICVGEGLARMELFLFTILQNFNLKSLIDPKDLDTTPVNGFASVPPFYQLCFIPV</p> <p>TRTRPLEQKLISEEDLAANDILDYKDDDDKV</p>
Tag:	C-Myc/DDK
Predicted MW:	55.8 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.



[View online »](#)

RefSeq: [NP_000760](#)

Locus ID: 1557

UniProt ID: [P33261](#)

RefSeq Size: 1473

Cytogenetics: 10q23.33

RefSeq ORF: 1470


Synonyms: CPCJ; CYP2C; CYPIIC17; CYPIIC19; P450C2C; P450IIC19

Summary: This gene encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This protein localizes to the endoplasmic reticulum and is known to metabolize many xenobiotics, including the anticonvulsive drug mephenytoin, omeprazole, diazepam and some barbiturates. Polymorphism within this gene is associated with variable ability to metabolize mephenytoin, known as the poor metabolizer and extensive metabolizer phenotypes. The gene is located within a cluster of cytochrome P450 genes on chromosome 10q24. [provided by RefSeq, Jul 2008]

Protein Families: Druggable Genome

Protein Pathways: Arachidonic acid metabolism, Drug metabolism - cytochrome P450, Linoleic acid metabolism, Metabolic pathways, Metabolism of xenobiotics by cytochrome P450, Retinol metabolism

Product images:



Coomassie blue staining of purified CYP2C19 protein (Cat# [TP311377]). The protein was produced from HEK293T cells transfected with CYP2C19 cDNA clone (Cat# [RC211377]) using MegaTran 2.0 (Cat# [TT210002]).