

Product datasheet for TP311472

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

RAB12 (NM_001025300) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human RAB12, member RAS oncogene family (RAB12), 20 μg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC211472 representing NM 001025300

or AA Sequence: Red=Cloning site Green=Tags(s)

MDPGAALQRRAGGGGLGAGSPALSGGQGRRRKQPPRPADFKLQVIIIGSRGVGKTSLMERFTDDTFCEA CKSTVGVDFKIKTVELRGKKIRLQIWDTAGQERFNSITSAYYRSAKGIILVYDITKKETFDDLPKWMKMI DKYASEDAELLLVGNKLDCETDREITRQQGEKFAQQITGMRFCEASAKDNFNVDEIFLKLVDDILKKMPL

DILRNELSNSILSLQPEPEIPPELPPPRPHVRCC

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK
Predicted MW: 27.1 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 001020471

Locus ID: 201475 **UniProt ID:** Q6IQ22





RefSeq Size: 2138

Cytogenetics: 18p11.22


RefSeq ORF: 732

Summary: The small GTPases Rab are key regulators of intracellular membrane trafficking, from the

formation of transport vesicles to their fusion with membranes. Rabs cycle between an inactive GDP-bound form and an active GTP-bound form that is able to recruit to membranes different set of downstream effectors directly responsible for vesicle formation, movement, tethering and fusion. That Rab may play a role in protein transport from recycling endosomes to lysosomes regulating, for instance, the degradation of the transferrin receptor. Involved in

autophagy (By similarity).[UniProtKB/Swiss-Prot Function]

Product images:



Coomassie blue staining of purified RAB12 protein (Cat# TP311472). The protein was produced from HEK293T cells transfected with RAB12 cDNA clone (Cat# [RC211472]) using MegaTran 2.0 (Cat# [TT210002]).