

## Product datasheet for TP311582L

### BTK (NM\_000061) Human Recombinant Protein

#### Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human Bruton agammaglobulinemia tyrosine kinase (BTK), 1 mg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC211582 representing NM_000061 Red=Cloning site Green=Tags(s)

MAAVILESIFLKRSQQKKKTSPLNFKKRLFLLTVHKLSYYEYDFERGRRGSKKGSIDVEKITCVETVPE  
KNPPPERQIPRRGEESSEMEQISIIERFPYPFQVYDEGPLYVFSPTTEELRKRWIHQKLN VIRYNSDLVQ  
KYHPCFWIDGQYLCCSQTAKNAMGCQILENRNGSLKPGSSHRKTKKPLPPTPEEDQILKKPLPEPAAAP  
VSTSELKKVVALYDYMPMNANDLQLRKGDEYFILEESNLPWWRRARDKNGQEGYIPSNYVTEAEDSIEMYE  
WYSKHMTRSQAQELLKQEGKEGGFIVRDSKAGKYTVSVFAKSTGDPQGVIRHYVVCSTPQSQYYLAEKH  
LFSTIPELINYHQHNSAGLISRLKYPVSQQNKNAPSTAGLGYGSWEIDPKDLTFLKELGTGQFGVWKYK  
WRGQYDVAIKMIKEGSMSEDEFIEEAKVMMNLSHEKLVQLYGVCTKQRPIFIITEYMANGCLLNLYREMR  
HRFQTQQLLEMCKDVCEAMEYLESKQFLHRDLAARNCLVNDQGVVKSDFGLSRYVLDDDEYTSSVGSKFP  
VRWSPPEVLMYSKFSSKSDIWAFGVLMWEIYSLGKMPYERFTNSETAEHIAQGLRLYRPHLASEKVYTIM  
YSCWHEKADERPTFKILLSNILDVMDEES

TRTRPLEQKLISEEDLAANDILDYKDDDDKV


Tag:	C-Myc/DDK
Predicted MW:	76.1 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.



[View online »](#)

<b>Stability:</b>	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
<b>RefSeq:</b>	<a href="#">NP_000052</a>
<b>Locus ID:</b>	695
<b>UniProt ID:</b>	<a href="#">Q06187</a> , <a href="#">Q5JY90</a>
<b>RefSeq Size:</b>	2591
<b>Cytogenetics:</b>	Xq22.1
<b>RefSeq ORF:</b>	1977
<b>Synonyms:</b>	AGMX1; AT; ATK; BPK; IGHD3; IMD1; PSCTK1; XLA
<b>Summary:</b>	The protein encoded by this gene plays a crucial role in B-cell development. Mutations in this gene cause X-linked agammaglobulinemia type 1, which is an immunodeficiency characterized by the failure to produce mature B lymphocytes, and associated with a failure of Ig heavy chain rearrangement. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Dec 2013]
<b>Protein Families:</b>	Druggable Genome, Protein Kinase
<b>Protein Pathways:</b>	B cell receptor signaling pathway, Fc epsilon RI signaling pathway, Primary immunodeficiency

### Product images:



Coomassie blue staining of purified BTK protein (Cat# [TP311582]). The protein was produced from HEK293T cells transfected with BTK cDNA clone (Cat# [RC211582]) using MegaTran 2.0 (Cat# [TT210002]).