

# **Product datasheet for TP311922**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Inhibin beta C chain (INHBC) (NM\_005538) Human Recombinant Protein

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Recombinant protein of human inhibin, beta C (INHBC), 20 μg

Species: Human
Expression Host: HEK293T

**Expression cDNA Clone** >RC211922 representing NM\_005538 **or AA Sequence:** Red=Cloning site Green=Tags(s)

MTSSLLLAFLLLAPTTVATPRAGGQCPACGGPTLELESQRELLLDLAKRSILDKLHLTQRPTLNRPVSRA ALRTALQHLHGVPQGALLEDNREQECEIISFAETGLSTINQTRLDFHFSSDRTAGDREVQQASLMFFVQL PSNTTWTLKVRVLVLGPHNTNLTLATQYLLEVDASGWHQLPLGPEAQAACSQGHLTLELVLEGQVAQSSV ILGGAAHRPFVAARVRVGGKHQIHRRGIDCQGGSRMCCRQEFFVDFREIGWHDWIIQPEGYAMNFCIGQC PLHIAGMPGIAASFHTAVLNLLKANTAAGTTGGGSCCVPTAQRPLSLLYYDRDSNIVKTDIPDMVVEACG

CS

36.3 kDa

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Predicted MW:

Concentration: >0.05 µg/µL as determined by microplate BCA method

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

**Preparation:** Recombinant protein was captured through anti-DDK affinity column followed by conventional

chromatography steps.

**Note:** For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

**Stability:** Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 005529

**Locus ID:** 3626



#### Inhibin beta C chain (INHBC) (NM\_005538) Human Recombinant Protein - TP311922

**UniProt ID:** P55103

RefSeq Size: 1990

Cytogenetics: 12q13.3

1056 RefSeq ORF: Synonyms: **IHBC** 

**Summary:** This gene encodes a member of the TGF-beta (transforming growth factor-beta) superfamily of


> proteins. The encoded preproprotein is proteolytically processed to generate a subunit of homodimeric and heterodimeric activin complexes. The heterodimeric complex may function in the inhibition of activin A signaling. Transgenic mice overexpressing this gene exhibit defects

in testis, liver and prostate. [provided by RefSeq, Aug 2016]

**Protein Families:** Druggable Genome, Secreted Protein

**Protein Pathways:** Cytokine-cytokine receptor interaction, TGF-beta signaling pathway

## **Product images:**



Coomassie blue staining of purified INHBC protein (Cat# TP311922). The protein was produced from HEK293T cells transfected with INHBC cDNA clone (Cat# [RC211922]) using MegaTran 2.0 (Cat# [TT210002]).