

Product datasheet for TP312288L

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PIP5K3 (PIKFYVE) (NM_152671) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human phosphatidylinositol-3-phosphate/phosphatidylinositol 5-

kinase, type III (PIP5K3), transcript variant 3, 1 mg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC212288 representing NM_152671 or AA Sequence: Red=Cloning site Green=Tags(s)

MATDDKTSPTLDSANDLPRSPTSPSHLTHFKPLTPDQDEPPFKSAYSSFVNLFRFNKERAEGGQGEQQPL SGSWTSPQLPSRTQSVRSPTPYKKQLNEELQRRSSALGDLRACTYCRKIALSYAHSTDSNSIGEDLNALS DSACSVSVLDPSEPRTPVGSRKASRNIFLEDDLAWQSLIHPDSSNTPLSTRLVSVQEDAGKSPARNRSAS ITNLSLDRSGSPMVPSYETSVSPQANRTYVRTETTEDERKILLDSVQLKDLWKKICHHSSGMEFQDHRYW LRTHPNCIVGKELVNWLIRNGHIATRAQAIAIGQAMVDGRWLDCVSHHDQLFRDEYALYRPLQSTEFSET PSPDSDSVNSVEGHSEPSWFKDIKFDDSDTEQIAEEGDDNLANSASPSKRTSVSSFQSTVDSDSAASISL

NVELDNVNFHIKKPSKYPHVPPHPADQKGRR

TRTRPLEQKLISEEDLAANDILDYKDDDDK**V**

Tag: C-Myc/DDK

Predicted MW: 50 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.





PIP5K3 (PIKFYVE) (NM_152671) Human Recombinant Protein - TP312288L

RefSeq: NP 689884

 Locus ID:
 200576

 UniProt ID:
 Q9Y2I7

 RefSeq Size:
 1661

 Cytogenetics:
 2q34

 RefSeq ORF:
 1353

Synonyms: CFD; FAB1; HEL37; PIP5K; PIP5K3; ZFYVE29

Summary: Phosphorylated derivatives of phosphatidylinositol (Ptdlns) regulate cytoskeletal functions,

membrane trafficking, and receptor signaling by recruiting protein complexes to cell- and endosomal-membranes. Humans have multiple PtdIns proteins that differ by the degree and position of phosphorylation of the inositol ring. This gene encodes an enzyme (PIKfyve; also known as phosphatidylinositol-3-phosphate 5-kinase type III or PIPKIII) that phosphorylates the D-5 position in PtdIns and phosphatidylinositol-3-phosphate (PtdIns3P) to make PtdIns5P and PtdIns(3,5)biphosphate. The D-5 position also can be phosphorylated by type I PtdIns4P-5-kinases (PIP5Ks) that are encoded by distinct genes and preferentially phosphorylate D-4 phosphorylated PtdIns. In contrast, PIKfyve preferentially phosphorylates D-3 phosphorylated PtdIns. In addition to being a lipid kinase, PIKfyve also has protein kinase activity. PIKfyve regulates endomembrane homeostasis and plays a role in the biogenesis of endosome carrier vesicles from early endosomes. Mutations in this gene cause corneal fleck dystrophy (CFD); an autosomal dominant disorder characterized by numerous small white flecks present in all layers of the corneal stroma. Histologically, these flecks appear to be keratocytes distended with lipid and mucopolysaccharide filled intracytoplasmic vacuoles. Alternative splicing results in multiple transcript variants encoding distinct isoforms.[provided by RefSeq,

May 2010]


Protein Families: Druggable Genome

Protein Pathways: Endocytosis, Fc gamma R-mediated phagocytosis, Inositol phosphate metabolism, Metabolic

pathways, Phosphatidylinositol signaling system, Regulation of actin cytoskeleton



Product images:



Coomassie blue staining of purified PIKFYVE protein (Cat# [TP312288]). The protein was produced from HEK293T cells transfected with PIKFYVE cDNA clone (Cat# [RC212288]) using MegaTran 2.0 (Cat# [TT210002]).