

Product datasheet for **TP312394**

STAT3 (NM_213662) Human Recombinant Protein

Product data:


Product Type:	Recombinant Proteins
Description:	Recombinant protein of human signal transducer and activator of transcription 3 (acute-phase response factor) (STAT3), transcript variant 3, 20 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA	>RC212394 representing NM_213662
Clone or AA	Red=Cloning site Green=Tags(s)
Sequence:	<p>MAQWNQLQQLDTRYLEQLHQLYSDFSFMELRQFLAPWIESQDWAYAASKESHATLVFHNLLGEIDQQYSR FLQESNVLYQHNLRRRIKQFLQSRYLEKPMELARIVARCLWEESRLLQTAATAAQGGQANHPTAAVTEK QQMLEQHLQDVRKRVQDLEQKMKVVENLQDDDFNYKTLKSQGDMQDLNGNNQSVTRQKMQQLEQMLTAL DQMRRSIVSELAGLLSAMEYVQKTLTDEELADWKRRQQIACIGGPPNICLDRLNWTSLAESQLQTRQQ IKKLEELQQKVSYKGDPIVQHRPMLERIVELFRNLMKSAFVVERQPCMPMHPDRPLVIKTGVQFTTKVR LLVKFPELNYQLKIKVCIDKDSGDVAALRGSRKFNLGTNTKVMNMEESNNGSLSAEFKHLTLREQRCGN GGRANCDASLIVTEELHLITFETEVYHQGLKIDLETHSLPVVISNICQMPNAWASILWYNMLTNNPKNV NFFTTPPIGTWDQVAEVLWSQFSSTTKRGLSIEQLTTLAELKLLGPGVNYSGCQITWAKFCKENMAGKGF FWWLDNIIDLKYLALWNEGYIMGFISKERERAILSTKPPGTFLRFSESSKEGGVTFTWVEKDISG KTQIQSVEPYTKQQLNNMSFAEIIMGYKIMDATNILVSPLVLYPDIPKEEAFGKYCRPESQEHPADPG SAAPYLKTKFICVTPFIDAVWK</p> <p>TRTRPLEQKLISEEDLAANDILDYKDDDDKV</p>
Tag:	C-Myc/DDK
Predicted MW:	82.9 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.



[View online »](#)

Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_998827
Locus ID:	6774
UniProt ID:	P40763
RefSeq Size:	4819
Cytogenetics:	17q21.2
RefSeq ORF:	2166
Synonyms:	ADMIO; ADMIO1; APRF; HIES
Summary:	The protein encoded by this gene is a member of the STAT protein family. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein is activated through phosphorylation in response to various cytokines and growth factors including IFNs, EGF, IL5, IL6, HGF, LIF and BMP2. This protein mediates the expression of a variety of genes in response to cell stimuli, and thus plays a key role in many cellular processes such as cell growth and apoptosis. The small GTPase Rac1 has been shown to bind and regulate the activity of this protein. PIAS3 protein is a specific inhibitor of this protein. This gene also plays a role in regulating host response to viral and bacterial infections. Mutations in this gene are associated with infantile-onset multisystem autoimmune disease and hyper-immunoglobulin E syndrome. [provided by RefSeq, Aug 2020]
Protein Families:	Druggable Genome, Transcription Factors
Protein Pathways:	Acute myeloid leukemia, Adipocytokine signaling pathway, Chemokine signaling pathway, Jak-STAT signaling pathway, Pancreatic cancer, Pathways in cancer

Product images:



Coomassie blue staining of purified STAT3 protein (Cat# TP312394). The protein was produced from HEK293T cells transfected with STAT3 cDNA clone (Cat# [RC212394]) using MegaTran 2.0 (Cat# [TT210002]).