

Product datasheet for TP312425

AMHR2 (NM_020547) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins
Description: Recombinant protein of human anti-Mullerian hormone receptor, type II (AMHR2), 20 µg
Species: Human
Expression Host: HEK293T
Expression cDNA Clone or AA Sequence: >RC212425 representing NM_020547
Red=Cloning site **Green**=Tags(s)

MLGSLGLWALLPTAVEAPPNRRTCFFVFEAPGVRGSTKTLGELLDTGTLPRAIRCLYSRCCFGIWNLTQD
RAQVEMQGCGRDSDPGCESLHCDPSRAHPSPGSTLFTCSCGTDFCNANYSHLPPPSPGTPGSQGPQAA
PGESIWMALVLLGLFLLLLLLLLLSIILALLQRKNYRVRGEPVPEPRPDSGRDWSVELQELPELCFSQQVI
REGGHAVVWAGQLQGKLVAIKAFPPRSVAQFQAERALYELPGLQHDHIVRFITASRGGPGRLLSGPLLVL
ELHPKGSLSCHYLTYQTSWSSLRMALSQAQLAFLHEERWQNGQNKPGIAHRDLSSQNVLIREDGSCAI
GDLGLALVLPGLTQPPAWTPTQPQGPAAIMEAGTQRYMAPELLDKTLDLQDWGMALRRADIYSLALLWE
ILSRCPDLRPDSSPPPFQLAYEAEELGNTPTSDELWALAVQERRRPYIPSTWRCFATDPDGLRELLEDWCW
ADPEARLTAECVQQRLAALHPQESHFPESCPRGCPPLCPEDCTSIPAPTILPCRQPQRSACHFSVQQGP
CSRNPQPACTLSPV

TRTRPLEQKLISEEDLAANDILDYKDDDDKV


Tag: C-Myc/DDK
Predicted MW: 62.6 kDa
Concentration: >0.05 µg/µL as determined by microplate BCA method
Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation: Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note: For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage: Store at -80°C.



[View online »](#)

Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_065434
Locus ID:	269
UniProt ID:	Q16671
RefSeq Size:	1855
Cytogenetics:	12q13.13
RefSeq ORF:	1722
Synonyms:	AMHR; MISR2; MISRII; MR11
Summary:	This gene encodes the receptor for the anti-Mullerian hormone (AMH) which, in addition to testosterone, results in male sex differentiation. AMH and testosterone are produced in the testes by different cells and have different effects. Testosterone promotes the development of male genitalia while the binding of AMH to the encoded receptor prevents the development of the mullerian ducts into uterus and Fallopian tubes. Mutations in this gene are associated with persistent Mullerian duct syndrome type II. Alternatively spliced transcript variants encoding different isoforms have been identified. [provided by RefSeq, Sep 2009]
Protein Families:	Druggable Genome, Protein Kinase, Transmembrane
Protein Pathways:	Cytokine-cytokine receptor interaction, TGF-beta signaling pathway

Product images:



Coomassie blue staining of purified AMHR2 protein (Cat# TP312425). The protein was produced from HEK293T cells transfected with AMHR2 cDNA clone (Cat# [RC212425]) using MegaTran 2.0 (Cat# [TT210002]).