

Product datasheet for TP312434L

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

ASAH1 (NM_004315) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human N-acylsphingosine amidohydrolase (acid ceramidase) 1

(ASAH1), transcript variant 2, 1 mg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone

or AA Sequence:

>RC212434 representing NM_004315 Red=Cloning site Green=Tags(s)

MNCCIGLGEKARGSHRASYPSLSALFTEASILGFGSFAVKAQWTEDCRKSTYPPSGPTYRGAVPWYTINL DLPPYKRWHELMLDKAPMLKVIVNSLKNMINTFVPSGKVMQVVDEKLPGLLGNFPGPFEEEMKGIAAVTD IPLGEIISFNIFYELFTICTSIVAEDKKGHLIHGRNMDFGVFLGWNINNDTWVITEQLKPLTVNLDFQRN NKTVFKASSFAGYVGMLTGFKPGLFSLTLNERFSINGGYLGILEWILGKKDAMWIGFLTRTVLENSTSYE EAKNLLTKTKILAPAYFILGGNQSGEGCVITRDRKESLDVYELDAKQGRWYVVQTNYDRWKHPFFLDDRR

TPAKMCLNRTSQENISFETMYDVLSTKPVLNKLTVYTTLIDVTKGQFETYLRDCPDPCIGW

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Predicted MW: 46.3 kDa

Concentration: $>0.05 \mu g/\mu L$ as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 004306





Locus ID: 427

UniProt ID: Q13510, Q53H01, A8K0B6

RefSeq Size: 2503 Cytogenetics: 8p22 RefSeq ORF: 1233

Synonyms: AC; ACDase; ASAH; PHP; PHP32; SMAPME

Summary: This gene encodes a member of the acid ceramidase family of proteins. Alternative splicing


results in multiple transcript variants, at least one of which encodes a preproprotein that is proteolytically processed. Processing of this preproprotein generates alpha and beta subunits that heterodimerize to form the mature lysosomal enzyme, which catalyzes the degradation of ceramide into sphingosine and free fatty acid. This enzyme is overexpressed in multiple human cancers and may play a role in cancer progression. Mutations in this gene are associated with the lysosomal storage disorder, Farber lipogranulomatosis, and a neuromuscular disorder, spinal muscular atrophy with progressive myoclonic epilepsy.

[provided by RefSeq, Oct 2015]

Protein Families: Druggable Genome

Protein Pathways: Lysosome, Metabolic pathways, Sphingolipid metabolism

Product images:



Coomassie blue staining of purified ASAH1 protein (Cat# [TP312434]). The protein was produced from HEK293T cells transfected with ASAH1 cDNA clone (Cat# [RC212434]) using MegaTran 2.0 (Cat# [TT210002]).