

## Product datasheet for **TP312592L**

### CDK8 (NM\_001260) Human Recombinant Protein

#### Product data:

**Product Type:** Recombinant Proteins

**Description:** Recombinant protein of human cyclin-dependent kinase 8 (CDK8), 1 mg

**Species:** Human

**Expression Host:** HEK293T

**Expression cDNA Clone** >RC212592 representing NM\_001260

**or AA Sequence:** **Red**=Cloning site **Green**=Tags(s)

MDYDFKVKLSSERERVEDLFYEGCKVGRGTYGHVYKAKRKDGKDDKDYALKQIEGTGISMSACREIALL  
RELKHPNVISLQKVFLSHADRKVVLLFDYAEHDLWHIIKFHRASKANKKPVQLPRGMVKSLLYQILDGIH  
YLHANWVLHRDLKPANILVMGEGPERGRVKIADMGFARLFNSPLKPLADLDPVVVTFWYRAPELLGARH  
YTKAIDIWAIGCIFAELLTSEPIFHCQRQEDIKTSNPYHHDQLDRIFNVMGFADKDWEDIKKMPEHSTLM  
KDFRRNTYTNCSLIKMEKHVKPDSKAFHLLQKLLTMDPIKRITSEQAMQDPYFLEDPLPTSDVFAFGCQ  
IPYPKREFLTEEPPDDKGDKNQQQQGNNHTNGTGHPGNQSSHTQGPPKKVRVVPPTTSSGGLIMITSD  
YQRSNPHAAYPNPGPSTSQPQSSMGYSATSQQPPQYSHQTHRY

**TRTRPLEQKLISEEDLAANDILDYKDDDDKV**

**Tag:** C-Myc/DDK

**Predicted MW:** 53.1 kDa

**Concentration:** >0.05 µg/µL as determined by microplate BCA method

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

**Preparation:** Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.

**Note:** For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.

**Storage:** Store at -80°C.

**Stability:** Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.

**RefSeq:** [NP\\_001251](#)



[View online »](#)

Locus ID: 1024

UniProt ID: [P49336](#)

RefSeq Size: 1772

Cytogenetics: 13q12.13


RefSeq ORF: 1389

Synonyms: IDDHBA; K35

**Summary:** This gene encodes a member of the cyclin-dependent protein kinase (CDK) family. CDK family members are known to be important regulators of cell cycle progression. This kinase and its regulatory subunit, cyclin C, are components of the Mediator transcriptional regulatory complex, involved in both transcriptional activation and repression by phosphorylation of the carboxy-terminal domain of the largest subunit of RNA polymerase II. This kinase regulates transcription by targeting the cyclin-dependent kinase 7 subunits of the general transcription initiation factor IIH, thus providing a link between the Mediator complex and the basal transcription machinery. Multiple pseudogenes of this gene have been identified. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Oct 2016]

**Protein Families:** Druggable Genome, Protein Kinase, Transcription Factors

### Product images:



Coomassie blue staining of purified CDK8 protein (Cat# [TP312592]). The protein was produced from HEK293T cells transfected with CDK8 cDNA clone (Cat# [RC212592]) using MegaTran 2.0 (Cat# [TT210002]).