

Product datasheet for TP312775M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

ICAM2 (NM_001099789) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Purified recombinant protein of Homo sapiens intercellular adhesion molecule 2 (ICAM2),

transcript variant 4, 100 µg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC212775 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MSSFGYRTLTVALFTLICCPGSDEKVFEVHVRPKKLAVEPKGSLEVNCSTTCNQPEVGGLETSLDKILLD EQAQWKHYLVSNISHDTVLQCHFTCSGKQESMNSNVSVYQPPRQVILTLQPTLVAVGKSFTIECRVPTVE PLDSLTLFLFRGNETLHYETFGKAAPAPQEATATFNSTADREDGHRNFSCLAVLDLMSRGGNIFHKHSAP KMLEIYEPVSDSQMVIIVTVVSVLLSLFVTSVLLCFIFGQHLRQQRMGTYGVRAAWRRLPQAFRP

Tag: C-Myc/DDK

Predicted MW: 28.3 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: <u>NP 001093259</u>

Locus ID: 3384





Synonyms:

UniProt ID: P13598, Q6FHE2

RefSeq Size: 1153 Cytogenetics: 17q23.3 825

RefSeq ORF:

Summary: The protein encoded by this gene is a member of the intercellular adhesion molecule (ICAM)

family. All ICAM proteins are type I transmembrane glycoproteins, contain 2-9

immunoglobulin-like C2-type domains, and bind to the leukocyte adhesion LFA-1 protein. This

protein may play a role in lymphocyte recirculation by blocking LFA-1-dependent cell

adhesion. It mediates adhesive interactions important for antigen-specific immune response,

NK-cell mediated clearance, lymphocyte recirculation, and other cellular interactions important for immune response and surveillance. Several transcript variants encoding the


same protein have been found for this gene. [provided by RefSeq, Jul 2008]

Protein Families: ES Cell Differentiation/IPS, Transmembrane

CD102

Protein Pathways: Cell adhesion molecules (CAMs), Natural killer cell mediated cytotoxicity

Product images:



Coomassie blue staining of purified ICAM2 protein (Cat# [TP312775]). The protein was produced from HEK293T cells transfected with ICAM2 cDNA clone (Cat# [RC212775]) using

MegaTran 2.0 (Cat# [TT210002]).