

Product datasheet for TP312815

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

NANOS2 (NM 001029861) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human nanos homolog 2 (Drosophila) (NANOS2), 20 µg

Species: Human HEK293T **Expression Host:**

Expression cDNA >RC212815 protein sequence Clone or AA Sequence: Red=Cloning site Green=Tags(s)

> MQLPPFDMWKDYFNLSQVVWALIASRGQRLETQEIEEPSPGPPLGQDQGLGAPGANGGLGTLCNFCKHNG ESRHVYSSHQLKTPDGVVVCPILRHYVCPVCGATGDQAHTLKYCPLNGGQQSLYRRSGRNSAGRRVKR

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

C-Myc/DDK Tag:

Predicted MW: 15 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

> 80% as determined by SDS-PAGE and Coomassie blue staining **Purity:**

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Recombinant protein was captured through anti-DDK affinity column followed by conventional Preparation:

chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Store at -80°C. Storage:

Stability: Stable for 12 months from the date of receipt of the product under proper storage and handling

conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 001025032

Locus ID: 339345 **UniProt ID:** P60321 RefSeq Size: 1577 Cytogenetics: 19q13.32





RefSeq ORF: 414

Synonyms: NOS2; ZC2HC12B


Summary: Plays a key role in the sexual differentiation of germ cells by promoting the male fate but

suppressing the female fate. Represses the female fate pathways by suppressing meiosis, which in turn results in the promotion of the male fate. Maintains the suppression of meiosis by preventing STRA8 expression, which is required for premeiotic DNA replication, after CYP26B1 is

decreased. Regulates the localization of the CCR4-NOT deadenylation complex to P-bodies and plays a role in recruiting the complex to trigger the degradation of mRNAs involved in meiosis. Required for the maintenance of the spermatogonial stem cell population. Not essential for the assembly of P-bodies but is required for the maintenance of their normal state (By similarity).

[UniProtKB/Swiss-Prot Function]

Product images:



Coomassie blue staining of purified NANOS2 protein (Cat# TP312815). The protein was produced from HEK293T cells transfected with NANOS2 cDNA clone (Cat# [RC212815]) using MegaTran 2.0 (Cat# [TT210002]).